

Neutrino Flux Prediction for the NuMI Beamline

Leonidas Aliaga Soplin

Lima, Peru

Master of Science, College of William and Mary, 2011

Master of Physics, PUCP, 2008

Bachelor of Physics, PUCP, 2002

A Dissertation presented to the Graduate Faculty
of the College of William and Mary in Candidacy for the Degree of
Doctor of Philosophy

Department of Physics

The College of William and Mary
March 2016

©2016
Leonidas Aliaga Soplin
All rights reserved.

TABLE OF CONTENTS

List of Tables	vi
List of Figures	viii
CHAPTER	
1 Introduction	2
1.1 Importance of the neutrino flux determination	2
2 NuMI beamline	10
2.1 Conventional neutrino beams	10
2.2 NuMI description	15
2.2.1 Primary proton beam	15
2.2.2 Target	17
2.2.3 Magnetic horns	19
2.2.4 Decay pipe and Absorbers	25
2.2.5 NuMI simulation	27
2.3 NuMI fluxes	30
2.3.1 Beam components	35
2.4 Hadronic cascade in NuMI	37
2.4.1 Neutrino ancestries	37
2.4.2 Interaction map	38
2.4.3 Hadronic production	42
2.4.4 Beam attenuation	42
2.5 Focusing uncertainties	44
2.5.1 Transverse horn offset	48

2.5.2	Baffle scraping	48
2.5.3	POT counting	49
2.5.4	Horn current	49
2.5.5	Horn inner conductor shape	49
2.5.6	Longitudinal target offset	51
2.5.7	Water layer uncertainty	52
3	Overview of neutrino flux determination strategies	55
3.1	Using external data	56
3.2	Using a multi-beam fitting	58
3.3	Low-nu method	61
3.4	Neutrino - electron scattering constraint	62
3.5	Minerva strategy	63
4	Correcting the simulation using hadron production data	65
4.1	Definitions	66
4.1.1	Total cross-section	66
4.1.2	Differential cross-section	67
4.2	Procedure	67
4.2.1	Attenuation Corrections	68
4.2.2	Particle Production	70
4.2.3	Extending data coverage	77
4.2.4	Procedure when there is no data	80
4.2.5	Uncertainty propagation	83
4.3	Data available	84
4.4	The thick target datasets	86
4.4.1	Thick target pion production	88
4.4.2	Thick target kaon production	97

4.5	The thin target hadron production datasets	104
4.5.1	NA49	104
4.5.2	Barton et al.	106
4.5.3	MIPP thin target	107
4.5.4	Thin target pion production	107
4.5.5	Thin target kaon production	112
4.5.6	Thin target nucleon production	116
4.6	Thin target data extension in Generation 2	124
4.6.1	Interactions of nucleons not covered by thin target data . .	127
4.6.2	Incident mesons and other interactions	127
4.7	Thin target inelastic and absorption cross section datasets	129
4.7.1	Proton - carbon inelastic cross-section	129
4.7.2	π^\pm absorption cross-sections	129
4.7.3	K^\pm absorption cross-sections	130
4.7.4	Technical implementation of the absorption and inelastic cross section	131
4.8	PPFX	137
4.8.1	MakeReweight	139
4.8.2	Data and MC Inputs	140
4.8.3	InteractionChainData	140
4.8.4	ReweightDriver	141
4.8.5	Reweighters	141
5	A priori MINERvA Low Energy Beam results	143
5.1	MINERvA flux Generation 0	144
5.2	MINERvA flux Generation 1	147
5.3	MINERvA flux Generation 2	149

5.3.1	Results	158
5.4	Discussion	171
6	Additional flux constraining using MINERvA in-situ measurements . .	174
6.1	Low-nu flux at MINERvA	175
6.1.1	Generation 2 vs low-nu flux comparison	179
6.2	Additional constraint using neutrino electron scattering events . .	183
6.3	Comparison of Gen2-thick with Generation 0 and Generation 1 .	186
7	Medium Energy Beam results	196
7.1	Gen2-thin flux for ME MINERvA	197
7.2	Gen2-thin for Nova ND	203
8	Conclusions	208
APPENDIX A		
	Supplementary material for chapter 2	210
A.1	Focusing components	210
A.2	Flux HP leading to neutrino in NOvA	218
APPENDIX B		
	NuMI flux tables	226
	Bibliography	231

LIST OF TABLES

2.1	Main decay modes into neutrinos	11
2.2	LE and ME target parameters	19
2.3	LE NuMI runs	31
2.4	Relevant volumes for hadron production correction	44
2.5	MINERvA LE flux playlists	51
4.1	Material scaling uncertainties	80
4.2	NA49 systematics uncertainties	106
4.3	Meson absorption cross-section uncertainty implemented in Generation137	
4.4	PPFX reweighters	142
5.1	Flux used in MINERvA publications	149
5.2	Integrated flux for ν_μ and $\bar{\nu}_\mu$ in Gen2-thin and Gen2-thick	171
5.3	Disagreement between Gen2-thin and Gen2-thick for ν_μ	172
6.1	Low-nu bins	177
6.2	χ^2/NDF for Generation 2 and low-nu difference	182
6.3	χ^2/NDF for Gen2-thick and Gen2-thin difference	182
6.4	Integrated flux ratios between generations	188
7.1	Integrated fluxes for Gen2-thin	198

B.1	Generation 2 - thin LE flux results at MINERvA	227
B.2	Generation 2 - thick LE flux results at MINERvA	228
B.3	Generation 2 - thin ME flux results at MINERvA	229
B.4	Generation 2 - thin ME flux results at NOvA ND	230

LIST OF FIGURES

1.1	A two detector neutrino oscillation experiment	3
1.2	F/N MINOS flux	5
1.3	Charged inclusive neutrino cross-section per nucleon	7
1.4	Coherent π^\pm production cross-section	8
2.1	Energy relation between a neutrino and its meson parent	12
2.2	Diagram of a horn and a magnetic horn	13
2.3	NuMI beamline schematic view	15
2.4	Plan elevation view of NuMI Beam Facility	16
2.5	Photograph of the LE NuMI target	17
2.6	LE and ME target draws	18
2.7	Photograph of the Horn 1	20
2.8	Horn 1 magnetic field	22
2.9	Focusing components	24
2.10	Focusing components for LE010z185i	24
2.11	π^+ and $K^- \nu_\mu$ parent focusing component ratios for LE010z185i . . .	26
2.12	MC ν_μ fluxes for LE runs	32
2.13	π^+ production in the target	33
2.14	MC ν_μ fluxes for ME runs	35
2.15	Neutrino components for LE and ME beams	36
2.16	Neutrino ancestries at MINERvA	39

2.17	Interaction map for ν_μ in MINERvA	40
2.18	Incident protons for ν_μ in MINERvA	41
2.19	Hadron production in pC leading to ν_μ flux in MINERvA	43
2.20	Material traversed of particles leading to ν_μ flux in MINERvA	45
2.21	LE010z185i focusing fractional shifts	47
2.22	Effect of the new horn model	50
2.23	Longitudinal target position effect	52
2.24	Water layer draws	53
2.25	Water layer effect	54
3.1	Hadron production experiments for neutrino flux (HARP and NA61)	59
3.2	Minos beam fit results	60
3.3	neutrino - electron scattering Feynman diagram	62
4.1	Particle beam traversing a volume	69
4.2	The survival correction	69
4.3	Attenuation correction when an interaction happened ($\sigma = 193.6$ mb)	71
4.4	Attenuation correction when an interaction happened ($\sigma_{MC} = 368$ mb)	71
4.5	FLUKA energy scaling correction	72
4.6	Scaling NA49 to NA61	74
4.7	Fractional difference between NA49 and NA61 applying scaling	75
4.8	Skubic data fit	79
4.9	Example of geant4 model spread	81
4.10	Example of the multi-universe technique	84
4.11	Schematic view of MIPP detector	87

4.12	Interactions per π^+ that exits the target for p_Z in 0-20 GeV/c	89
4.13	Interactions per π^+ and K^+ that exit the target for p_Z in 20-80 GeV/c	90
4.14	MIPP NuMI π^+ and π^- data	91
4.15	MIPP NuMI π data statistical uncertainties	92
4.16	MIPP NuMI π data systematic uncertainties	93
4.17	Bin-to-bin correlation effect on MIPP NuMI systematic uncertainties	94
4.18	MIPP NuMI charged pions weights	96
4.19	MIPP NuMI yield ratios	97
4.20	MIPP NuMI K/π statistical uncertainties	98
4.21	MIPP NuMI K/π systematic uncertainties	100
4.22	MIPP NuMI charged kaons weights	102
4.23	MIPP NuMI neutral kaons weights	103
4.24	Schematic view of NA49 detector	105
4.25	NA49 charged pion production in proton - carbon interactions	108
4.26	Barton et al π^+ data in proton - carbon interactions	109
4.27	NA49 statistical uncertainty for pion production	111
4.28	Thin target weights for charged pions	113
4.29	NA49 charged kaon production in proton carbon interactions	114
4.30	MIPP pC yield ratios	115
4.31	NA49 statistical uncertainty for kaon production	117
4.32	MIPP pC statistical uncertainties for K/π	118
4.33	MIPP pC systematic uncertainties for K/π	119
4.34	Thin target weights for charged kaons	120
4.35	Thin target weights for neutral kaons	121

4.36	NA49 proton production in proton carbon interactions	123
4.37	NA49 neutron production in proton carbon interactions	124
4.38	NA49 nucleon statistical uncertainties	125
4.39	Thin target weights for nucleon production	126
4.40	Proton-carbon inelastic cross sections.	130
4.41	π C and π Al data and MC absorption cross-section	131
4.42	π data - MC absorption cross-section data - MC comparisons	132
4.43	Charged kaon on C and Al data and MC absorption cross-section . .	133
4.44	K^+ data - MC absorption cross-section data - MC comparisons . . .	134
4.45	K^- data - MC absorption cross-section data - MC comparisons . . .	135
4.46	PPFX flowchart.	138
5.1	Generation 0 flux	146
5.2	Generation 1 flux	148
5.3	Generation 0 and Generation 1 flux fractional uncertainties	150
5.4	Interactions per neutrino in Gen2-thin	154
5.5	Interactions per neutrino in Gen2-thick	155
5.6	Nucleon-A interaction per neutrino covered in Generation 2	157
5.7	NuMI material traversed	159
5.8	Gen2-thin ν_μ flux	161
5.9	Gen2-thick ν_μ flux	162
5.10	Gen2-thin $\bar{\nu}_\mu$ flux	163
5.11	Gen2-thick $\bar{\nu}_\mu$ flux	164
5.12	Gen2-thin ν_e flux	165
5.13	Gen2-thick ν_e flux	166

5.14	Fractional uncertainties for Gen2-thin	167
5.15	Fractional uncertainties for Gen2-thick	168
5.16	Energy bin to bin flux correlation	170
5.17	Flux integral correlation for Gen2-thin	171
5.18	Flux integral correlation for Gen2-thick	172
5.19	ν_μ Gen2-thin and Gen2-thick comparison	173
5.20	$\bar{\nu}_\mu$ Gen2-thin and Gen2-thick comparison	173
6.1	Overlap and ν cross-sections	176
6.2	ν_μ low-nu flux	178
6.3	Gen2-thin ν_μ flux vs low-nu	180
6.4	Gen2-thick ν_μ flux vs low-nu	181
6.5	Electron energy distribution from ν -e scattering	184
6.6	Effect of the νe constraint on the Generation 1 integrated flux	185
6.7	Effect of the νe constraint on the Gen2-thin flux	187
6.8	ν_μ flux Gen2-thin vs Generation 0	189
6.9	$\bar{\nu}_\mu$ flux Gen2-thin vs Generation 0	190
6.10	ν_μ flux Gen2-thin vs Generation 1	191
6.11	$\bar{\nu}_\mu$ flux Gen2-thin vs Generation 1	192
6.12	ν_e flux Gen2-thin vs Generation 1	193
6.13	ν_μ flux Gen2-thin vs Generation 1 + neutrino electron scattering constraint	194
6.14	ν_e flux Gen2-thin vs Generation 1 + neutrino electron scattering constraint	195
7.1	ME PPFX coverage for ν_μ in MINERvA	199

7.2	ME Gen2 - thin ν_μ flux at MINERvA	200
7.3	Fractional Uncertainties for ν_μ ME Gen2-thin flux in MINERvA. . .	201
7.4	Bin to bin correlation of ME Gen2-thin at MINERvA	202
7.5	ME Gen2-thin interactions and material coverages for ν_μ in NOvA .	204
7.6	Gen2-thin ME NOvA flux	205
7.7	Fractional Uncertainties for ν_μ ME Gen2 - thin flux in NOvA. . . .	206
7.8	Comparison between Gen2-thin and FLUGG ν_μ flux NOvA ND . .	207
A.1	Longitudinal vs. transverse momentum yield per π^+ ν_μ parent	211
A.2	Unfocused components of π^+ ν_μ parent	212
A.3	H2 only components of π^+ ν_μ parent	213
A.4	Horn1 only components of π^+ ν_μ parent	214
A.5	Underfocused components of π^+ ν_μ parent	215
A.6	Overfocused components of π^+ ν_μ parent	216
A.7	Other components of π^+ ν_μ parent	217
A.8	Focusing components for NOvA ND	219
A.9	Muon and electron neutrino parents in NOvA	220
A.10	Muon and electron neutrino grandparents in NOvA	221
A.11	Interaction map for ν_μ in NOvA	222
A.12	Incident protons for ν_μ in NOvA	223
A.13	Hadron production leading to ν_μ in NOvA	224
A.14	Beam attenuation for particles leading to ν_μ in NOvA	225

NEUTRINO FLUX PREDICTION FOR THE NUMI BEAMLINE

CHAPTER 1

Introduction

The determination of the neutrino flux in any conventional neutrino beam presents a challenge for the current and future short and long baseline neutrino experiments. The uncertainties associated with the production and attenuation of the hadrons in the beamline materials along with those associated with the beam optics have a big effect on the knowledge flux spectrum. For experiments like MINERvA, understanding the flux is crucial since it enters directly into every of neutrino-nucleus cross-section measurements.

1.1 Importance of the neutrino flux determination

Since the late 1990's, there has been overwhelming evidence that the three types of neutrinos, ν_e , ν_μ and ν_τ , oscillates when they travel through vacuum or matter. Since then, many experiments have been proposed and developed to measure the parameters associated with neutrino oscillations.

In many circumstances, considering only the oscillation between two neutrinos

is a good approximation. This is the case for neutrinos generated in the Earth atmosphere ($\nu_\mu \leftrightarrow \nu_\tau$) or, in the Sun or in conventional neutrino beams ($\nu_e \leftrightarrow \nu_\mu$). In these cases, the probability that a neutrino that has been produced in one flavor (ν_α) will be measured in another flavor (ν_β) is given by:

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \sin^2(2\theta) \sin^2 \left(\frac{1.27 \Delta m_{ij}^2 L}{E} \right) \quad (1.1)$$

Where L is the distance traveled by the neutrino and E , its energy. The factor “1.27” supposes that E is expressed in GeV, L in Km and the squared neutrino mass difference Δm^2 in eV^2/c^4 units.

Long baseline neutrino experiments use a conventional neutrino beam (described in chapter 2) and a pair of detectors, one close to the neutrino production point (the Near detector) and the other further away (the Far detector). The location of the Far detector and the beam energy spectrum are chosen to measure the oscillation maxima of equation 1.1. Having the two detectors share the same beam reduces systematic uncertainties on the neutrino flux, the cross-section model, and potentially also the detector performance. Figure 1.2 shows a sketch of this two detector strategy.




FIG. 1.1: A two detector neutrino oscillation experiment.

If the cross-section is well-understood, the oscillation probability can be calculated as:

$$P(E_i) = \left(\frac{N_2}{N_1} \right)_i \left(\frac{A_1}{A_2} \right)_i \left(\frac{\phi_1}{\phi_2} \right)_i \quad (1.2)$$

where $A_{1,2}$ represents the acceptance and efficiency of each detector and $\phi_{1,2}$ is the neutrino flux, and N_1 and N_2 are the events measured in a particular energy bin.

However, this is not simple. The flux uncertainty partially cancels since the Near Detector sees a distributed neutrino source and the Far Detector sees a point neutrino source (for instance, MINOS Far detector is ~ 1000 Km away). The cross-sections do not necessarily cancel if the detectors are not made of the same materials with the same response and acceptance.

As an example, figure 1.2 shows the ratio of the flux at the MINOS Far Detector divided by the flux at the MINOS Near Detector ($\frac{\phi_1}{\phi_2}$ in equation 1.2). The white boxes in the figure represent the prediction from the MINOS flux simulation. However, after MINOS tuned their simulation by constraining the hadron production in the target (see later section 3.2), the shape of the ratio changes. That means that knowing the flux and how well the underlying physics model is representing nature is absolutely relevant for the accuracy of the oscillation parameters.

On the other hand, the cross-sections are also a source of uncertainty for neutrino oscillation experiments. In particular, the 0.1-20 GeV region where many experiments are running is complicated because the neutrino - nucleus cross-sections are not well known, as can be seen in figures 1.3a and 1.3b. Some uncertainties and disagreements of these datasets could come from the uncertainties in the nuclear effects or from a poor knowledge of the neutrino flux.

Future experiments like DUNE will consist of liquid Argon (LAr) and it is crucial to understand the nuclear effect in the neutrino interactions models [2].




FIG. 1.2: MINOS Far over Near flux ratio (taken from [1]).

Large errors in cross section measurements and disagreements between experiments lead to systematic uncertainties in oscillation measurements.

The MINERvA experiment is dedicated to study ν_μ and $\bar{\nu}_\mu$ cross-sections neutrino in the 1-20 GeV range with high statistics for inclusive and exclusive channels and with interactions on several nuclear targets (C, Pb, Fe, H₂O and He) to investigate nuclear effects.


MINERvA is located on-axis in the NuMI beamline in the MINOS Target Hall and it is designed to have a good reconstruction of the neutrino interactions [4]. It consists of different nuclear targets that are followed by a core of scintillator strips (called the tracker) and surrounded by electromagnetic and hadronic calorimeters. One of the main challenges comes from the uncertainty in the neutrino flux.

A neutrino cross-section is calculated according to the equation 1.3 as the reconstructed neutrino interactions divided by the neutrino flux and the number of target nuclei (T). Any uncertainty in the flux enters directly to the cross-section calculation. This is the reason why knowing the flux is crucial for MINERvA.


$$\sigma(E) = \frac{N(E)}{\phi(E) \times T} \quad (1.3)$$

As an example of this challenge, figure 1.4 shows the coherent charged pion production cross-section in ν_μ interactions recently published by MINERvA [5]. These are the first high statistics measurements made of this channel. The uncertainties are dominated by the flux, limiting the precision of the measurement (see figure 1.4b). This result is very important for the neutrino physics community since the neutral current analog of this channel is a potential source of background in oscillation experiments.

This thesis is about determining the NuMI flux for the MINERvA experiment developing a procedure and tools that can be used to determine the flux for any



(a) ν_μ



(b) $\bar{\nu}_\mu$

FIG. 1.3: ν_μ and $\bar{\nu}_\mu$ per nucleon charged current cross-sections [3].





FIG. 1.4: Coherent production of π^\pm in ν_μ [5].

detector that sees neutrinos from NuMI. The structure is the following: after a description of the NuMI geometry and capabilities (chapter 2), we are going to develop our multi-layer approach described in chapter 3 consisting in the procedure explained in chapter 4. In chapters 5 - 7 we will present and analyze our results.

CHAPTER 2

NuMI beamline

I start this chapter introducing the aspects of the design and construction of conventional neutrino beams as a general context to describe the NuMI beamline. Not all parts of NuMI are included here, just the relevant components for understanding its geometry and capabilities in the context of this thesis. The major challenges that are presented in detail are the focusing system and the hadronic cascade from the primary proton beam to the weak decay producing a neutrino in the beamline. All plots are based on direct output of the simulation, i.e., no correction is applied to the hadronic model. The MINERvA strategy to correct the hadron production mismodeling is presented in the chapter 3.

2.1 Conventional neutrino beams

The basic concept of conventional neutrino beams is to extract a very intense proton beam from an accelerator and collide it with a target to produce short lived particles, like pions and kaons, that will eventually decay to neutrinos. The targets are thick (a few interactions length long) but narrow enough to allow the produced particles to escape without interacting too many times. The particles produced have

a boost in the direction of the beamline but in order to enhance the neutrino beam, one or more Van der Meer's magnetic horns ([6], [7]) are placed to focus the mesons. After the horns, the particles enter a long vacuum or low density pipe where most of mesons decay. The remaining muon and hadrons beams are removed by absorbers.

The main decay modes that produce neutrinos are presented in table 2.1. Charged pions and charged kaons create predominantly muon neutrinos and a small component of electron neutrinos. Muons can also be generated in these decays and contribute to the electron neutrino flux but these are highly suppressed because they are mostly absorbed before they decay.

Decay	Chanel	Branching ratio (%)
1	$\pi^\pm \rightarrow \mu^\pm + \nu_\mu(\bar{\nu}_\mu)$	99.9877
2	$\pi^\pm \rightarrow e^\pm + \nu_e(\bar{\nu}_e)$	0.0123
3	$K^\pm \rightarrow \mu^\pm + \nu_\mu(\bar{\nu}_\mu)$	63.55
4	$K^\pm \rightarrow \pi^0 + e^\pm + \nu_e(\bar{\nu}_e)$	5.07
5	$K^\pm \rightarrow \pi^0 + \mu^\pm + \nu_\mu(\bar{\nu}_\mu)$	3.353
6	$K_L^0 \rightarrow \pi^\pm + e^\mp + \nu_e$	40.55
7	$K_L^0 \rightarrow \pi^\pm + \mu^\mp + \nu_\mu$	27.04
8	$\mu^\pm \rightarrow e^\pm + \nu_e(\bar{\nu}_e) + \bar{\nu}_\mu(\nu_\mu)$	100.0

TABLE 2.1: Main decay modes and their branching ratios of charged pion, charged kaons, neutral kaons and muons to neutrinos.

Considering a two body decay (like in decay types 1-3 at the table 2.1), we can calculate the energy of the neutrino, assuming that they are moving in a near forward direction, as a function of its meson parents (equation 2.1). Muon and meson masses are represented by m_μ and M , respectively, the angle of the neutrino flux respect to the pion parent momentum is θ_ν and the Lorentz factor ($\gamma = E_{\pi(K)}/M$):

$$E_\nu \approx \frac{(1 - \frac{m_\mu^2}{M^2})E_{\pi(K)}}{1 + \gamma^2 \tan^2 \theta_\nu} \quad (2.1)$$

For detectors located on the beam axis (on-axis), like MINERvA and MINOS, the angle θ_ν is zero. Then the energies of the neutrino and its parent have a linear

relation, $E_\nu \approx 0.43E_\pi$ and $E_\nu \approx 0.95E_K$ for pion and kaon parents, respectively. For off-axis experiments like NOvA, where $\theta_\nu \approx 12.26$ mrad, the relation becomes non-linear due to the energy dependency of the Lorentz factor. Figure 2.2 shows the functional form of the equation 2.1 for pion neutrino parents at three positions: on-axis detectors (black line), the center of the front face of NOvA Near detector (red line) and the center of MicroBooNE (in purple). For practical reasons, the horizontal axis is cut at 10 GeV, but all off-axis curves reach a maximum and then decrease their value monotonically.




FIG. 2.1: Functional form given in equation 2.1 for three positions in NuMI beamline: on-axis (MINERvA and MINOS) in black, off-axis (NOvA and MicroBooNE).

Some criteria when designing a conventional neutrino beam are the following:

- **The proton beam.** Depending of the physics goals, selecting the energy and intensity of the primary proton beam follow two rules of thumb: firstly, higher energy proton beams produce higher energy mesons and then more energetic neutrino fluxes. Secondly, pion and kaon multiplicities are approximately pro-

- **Target.** Increasing the target length has the benefit to generate more interactions of the primary protons, however, it can bring two undesirable effects: it enlarges the scattering of the produced mesons and, in conjunction with an intense proton beam, generates more interactions and a hotter target.

[illegible]

- **Focusing system.** The challenge in focusing is to cancel the transverse momentum (p_T) of as many hadrons that exit the target as possible ¹. This means that

13

higher p_T hadrons have to travel more distance in the magnetic horns and, at the same time, the interaction probability in the horns has to be minimized. Figure 2.2 displays the particle trajectories of pions entering to a horn. In this case, π^+ are focused and π^- are deflected due to direction of the magnetic field.

Modern magnetic horns are composed of 2-layered thin coaxial sheet conductor with a volume between the layers filled with low density gas (air, helium or argon). The current circulates in one direction in the inner layer and in the opposite direction in the outer layer generating a toroidal magnetic field between layers that falls as $1/R$ by Amperes law and has zero value outside the horn. Depending of the direction of the current (and the magnetic field) the horns focus charged particles with a particular sign and deflect particles with the opposite sign. In this way, the wanted neutrino helicity is enhanced while the undesired neutrino chirality (background) is minimized.

The sketch in the left lower part of figure 2.7 shows this characteristic in the context of NuMI description in the the next section. Furthermore, having a parabolic front face makes the total distance travel by the particle in the horn (and the total momentum change) proportional to p_T .

- **Decay Pipe.** Increasing the length of the decay pipe allows more and also higher pions to decay but makes fewer muons reach the absorbers and the number of electron neutrinos increases. The final length depends of the required neutrino energy and how much ν_e contamination is tolerable according the the physics goals of the experiment. Ideally, the pipes should be evacuated to eliminate any interactions but this requires more material in the windows of the pipe to keep the vacuum (there are always some concerns of catastrophic failures in the system). Pipes filled with helium or air reduces the neutrino spectrum by a few percentage but are good alternatives.

2.2 NuMI description

The NuMI design was motivated by neutrino oscillation studies ([8], [9]). The parameters of the main components have been optimized to meet the objectives of MINOS for the Low Energy (LE) run and NOvA for the Medium Energy (ME) run. Its high intensity neutrino flux has been used by other experiments like MINERvA to study cross-sections.

The NuMI neutrino beamline starts at Fermilab, Illinois, and travels through the crust of the Earth towards the Soudan Mine in Minnesota and then exits the Earth. In its travel, ~ 1 km from the starting point, it encounters the MINOS Near Detector and MINERvA, and at ~ 730 km the MINOS Far Detector.

A schematic view of the NuMI beamline is shown in figure 2.3. A detailed description of every part can be found in [8]. In this section, I will be focused on those components related to this thesis: the primary beam, target, magnetic horns and the decay pipe.




FIG. 2.3: Schematic view of NuMI beamline (taken from [8]).

2.2.1 Primary proton beam

The proton accelerator at Fermilab starts with H^- ions being accelerated to 400 MeV kinetic energy in the LINAC (**L**INear **A**Ccelerator [10]). Those ions

are converted to H^+ (protons) when they pass by a carbon foil and injected to the Booster synchrotron ([11]), where the protons are accelerated to 8 GeV kinetic energy. Some of these protons are used for the short baseline neutrino experiments before entering to the Main Injector (MI).

The MI ring is a synchrotron accelerator of 3.3 km circumference that, in turn, accelerates the protons to 120 GeV, which are then used for many purposes. It served as the proton (and antiproton²) source to the Tevatron, provides beamlines for fixed target experiments and test beams, it is also used to create neutrinos. Protons from NuMI are extracted in 6 batches³ in approximately 10 μ s in a “single turn”⁴ and bent 58 mrad downward towards the MINOS Near detector. Figure 2.4 shows a plan and elevation view of the NuMI Beam Facility. The beam is directed to the Target Hall, 41 m underground, that contains the target and the magnetic horns.




FIG. 2.4: Plan elevation view of NuMI Beam Facility (taken from [8]).

The beam spot size at the target is gaussian with $\sigma_x = 1.1$ mm and $\sigma_y = 1.2$ mm.

²The Main Injector provided protons to the antiproton accumulator up to 2011 when Tevatron ceased operations.

³During the Tevatron era, 5 batches were destined to NuMI and 1 for the antiproton accumulator for Tevatron and the NuMI extraction time was 8 μ s.

⁴“Single turn” technique is a fast deliver of the entire beam in a single extraction[12].

In the LE run, the cycle time was between 2.1 s and 2.4 s per spill with 2.2E13 POT/spill in 2005 rising to 3.6E13 POT/spill in 2012. The average beam power was ~ 250 kW. For ME run, an upgrade is currently in progress to reach 700 kW beam power for NOvA physics goals. At the time of writing, the power is already ~ 400 kW with a 1.3 s cycle time. The beam spot is wider than in LE with $\sigma_{x(y)} = 1.3$ mm RMS.

2.2.2 Target

The target is a rectangular graphite rod with 1.78 g/cm^3 density and it is segmented in rectangular pieces (called “fins”) rounded at the edges and stack along the beam direction with a small space between them. The target is cooled by water that circulates in a pipe bonded to the upper and lower edge of each fin. The whole target is enclosed in a helium container. Budal monitors are placed upstream to check the position of the target by scanning the beam in the horizontal or vertical direction ⁵.

A picture of the LE target can be seen in figure 2.5.



FIG. 2.5: Photograph of the LE NuMI target [14].

A sketch is shown in figure 2.6 for details aid. At the top, a longitudinal view of a couple of fins where X, Y and Z represent the distance between fins, the fin

⁵Budal Monitors measure the signal from electron and other particles that are kicked off when the proton beam interact in a fin. They provide a position dependent signal proportional to the beam intensity. The original proposal can be found in [13].

length and the length of the rounded side. At the bottom, a cross-sectional view of the LE (left) and ME (right) targets and their dimensions. The blue circle indicates the incident point of the proton beam.




FIG. 2.6: LE and ME target draws.

For LE run, the target consists of 47 vertical fins plus 1 additional horizontal fin with a Budal monitor ⁶. Each fin is 20 mm long (Y), 15 mm tall and 6.4 mm wide. The space between fins is 0.3 mm (X). The beam is center exactly in the center of the transverse view of the target. The Budal monitor is placed 167.2 mm upstream of the target for horizontal scanning.

For the ME run, the target consist of 48 vertical fins plus 2 additional fins with Budal monitors, one vertical and the other horizontal. Due to the higher beam

⁶The system fin with a Budal monitor is simply called “Budal Monitor” (BM) in the context of this thesis, unless otherwise indicated.

power, ME fins are larger than the LE ones. Each fin is 24 mm long (Y), 63 mm tall and 7.4 mm wide. The separation between fins is 0.5 mm (X). The location of the beam is at the center of the horizontal side but 3.7 mm from the top of the target. The Budal monitors have the same dimensions as the fins: the vertical one, for vertical scanning, is located 28.5 mm from the target, while the horizontal one, for horizontal scanning, is placed 57.5 mm from the target.

A 1.5 m long baffle protects the target and the horns (especially the neck of the horn) from any missteering of the beam and it helps monitor the beam. It consists of a 57 mm diameter graphite core with a 11 mm diameter hole, all encased in an aluminum tube and two 0.5 mm thickness beryllium windows. The target, Budal monitors and Baffle are mounted in the same carrier. This allows them to be moved together and control the distance respect to the horn when the beam configuration changes.

Table 2.2 summarizes the main differences between the target in LE and ME configurations.

	LE	ME
Cross Sectional view	6.4 x 15 mm ²	7.4 x 63 mm ²
Segment length	20 mm	24 mm
“Fins”	47 + 1 BMs	48 + 2 BMs
Total Length	960 mm ($\sim 2 \lambda$)	1200 mm ($\sim 2.5 \lambda$)

TABLE 2.2: Summary of LE and ME target parameters.

2.2.3 Magnetic horns

NuMI uses two magnetic horns, each one 3 m long. A current circulates through the inner conductor (IC) and returns by the outer conductor (OC) creating a toroidal magnetic field between them. The conductors are made of aluminum and the field part is filled with Argon. Figure 2.7 shows a not-to-scale sketch of the cross-sectional

view of Horn 1, indicating the direction of the current and the magnetic field affecting the particles that enter the horn from the target. The picture in figure 2.7 is a photograph of the real Horn 1 Inner Conductor in the welding machine before being placed in the Target Hall.




FIG. 2.7: Photograph of the Horn 1 in the welding machine before being place in the Target Hall [14]. In the bottom left corner, a sketch of the target and Horn 1 indicating some possible particle trajectories.

The IC surface is made of two parabolic surfaces of rotation, one upstream and the other downstream, welded together with a short cylindrical neck. The thicknesses of the parabolic parts are 2 mm and 3 mm for Horn 1 and Horn2 respectively, and the neck is 4.5 mm thick. The OC diameter of the Horn 1 is ~ 35 cm Horn 1 and for the Horn 2 is ~ 79 cm, with 25.4 mm thick.

The current pulsed is a half-sine wave with typical values 182.1 kA and 200 kA

for LE and ME respectively. The duration of the pulse is 2.3 ms and it is long enough to provide a stable magnetic field. The inside surfaces of the ICs are continuously sprayed with cooling water to remove heat.


A map of the magnetic field in the Horn 1 is shown in figure 2.8 a) with respect to the beamline direction (Z) and the vertical position (Y) in the simulation coordinate system. The NuMI simulation will be explained in section 2.2.5. Due to the transverse symmetry of the horns, the Y axis can be replaced by a radial axis. The dashed line represents the target in the nominal position for LE (more details about target longitudinal position will be given in section 2.3). The maximum magnetic field is around 2.7 T in the neck region, creating the need for a thicker conductor. The field decreases as $1/r$ as can be seen in the radial distribution in figure 2.8 b).

The relative distance between horns and the Horn 1 and the target are crucial to the focusing process. Horn 2 is located downstream from Horn 1 at 7 m and 16 m approximately in LE and ME respectively. Also, in the ME configuration, the target is pulled back from the Horn 1. The effect that this causes in changing the beam energy is explained in section 2.3.


One way to understand the effect of the focusing system is by splitting the neutrino flux spectrum in categories (called “focusing components”) with respect to how the neutrino parent travels from the target through the horns (as display in figure 2.9) and it depends of the absolute momentum and the relative value of the transverse momentum with respect to the longitudinal momentum. These categories are:

- **Unfocused:** mesons that pass through the necks of both horns (solid red line).

They have high momentum (typically more than 15 GeV/c) and very small p_T ($p_T < 0.1$ GeV).



(a) Longitudinal cross sectional distribution (the target in LE010z position configuration is drawn in dashed line as reference).



(b) Radial distribution.

FIG. 2.8: Horn1 magnetic field for 185i current configuration (182.1 kA).

- **Horn2-only:** mesons that pass through the neck of the first horn and cross the Horn 2 (dashed red line). They have an intermediate energy (typically between 9-15 GeV/c) and the transverse momentum creates a small deviation from the beamline.
- **Underfocused:** mesons that pass through both horns receiving a correction in Horn 1 and complemented by Horn 2 (solid orange line). The momentum of these particles in the LE mode, is in 5-15 GeV/c range and a $p_T > 0.2 \text{ GeV}/c$.
- **Horn1-only:** mesons that are affected only by the Horn 1 and not by the Horn 2. They have $p_T > 0.2 \text{ GeV}/c$ as in underfocused category but less momentum making the Horn 1 bending the particle to the Horn 2 neck. Mesons in this category used to decay into neutrino early.
- **Overfocused:** mesons with low momentum (typically less than 5 GeV in LE configuration) that are overcorrected by the Horn 1 and that are also corrected by the the Horn 2.

In the appendix A, figures A.2-A.7 show the kinematic distribution of π^+ neutrino parents when they exit the target per focusing components and per beam configuration.

Figure 2.10 shows the focusing components for the ν_μ flux at MINERvA in the LE101z185i beam configuration (the beam configuration convention is explained in the next section). In addition, the category “others” has been added for those neutrino parents born outside of the target (dashed black line). The total LE ν_μ flux that passes through MINERvA is shown as a solid black line.

The ratios of every focusing component with respect to the part of the ν_μ flux in MINERvA that has a π^+ parent and K^+ parent are shown in figure 2.15 a) and b), respectively. As will be seen later in section 2.4.1, most of the neutrinos with




FIG. 2.9: Focusing components.




FIG. 2.10: focusing components of ν_μ flux in LE010z185i that pass through MINERvA.

energy less than 20 GeV have a π^+ parent and greater than 20 GeV, a K^+ parent.

For π^+ we extend the neutrino energy to 20 GeV. As we expect, as we go from higher to lower momentum, unfocused, underfocused and overfocused pion decays populate the ν_μ flux. A narrow momentum range values allow the pions to be focused by just one horn. The very low energetic neutrinos come mostly from pions born outside of the target from secondary and tertiary hadrons.

The ratios for K^+ cover up to 40 GeV. The underfocused kaons become relevant around 5-20 GeV. For energies in 20-40 GeV, the horn2-only component kaons are predominant.

The NOvA Near Detector ν_μ flux spectrum and its focusing components are shown in figure A.8 in the appendix A.

2.2.4 Decay pipe and Absorbers

The Decay Pipe begins 46 m downstream of the NuMI Target Hall. It has a cylindrical shape with a 1 m inner radius, 675 m length, and is surrounded by a 9.5 mm thick wall made of iron. It is cooled by water and surrounded by poured concrete shielding. Almost all pions with energy less than 10 GeV decay to ν_μ in the Decay Pipe.

Originally the pipe was evacuated but later was filled with helium gas. The effect of adding He on the neutrino spectrum was of the order of 10% reduction around 3-4 GeV and 5% increase for higher energies. In the context of this thesis and motivated by the simulation name convention, the Decay Pipe inner volume is abbreviated as DVOL and its wall as DPIP.

Just downstream of the Decay Pipe a massive structure made of aluminum, steel and concrete is placed to absorb the residual hadrons still remaining in the cascade. Is is a box 5.5 m wide, 5.6 m tall, and 8.5 m long. Following the absorber,





FIG. 2.11: Beam focusing component ratios of ν_μ flux in the LE010z185i configuration that pass through MINERvA.

240 m of earth, mainly dolomite rock, separates the Absorber and the MINOS Hall. The rock shields experiments in the MINOS hall from the residual μ beam.

The NuMI components that were described in this section are relevant in the context of this thesis, as will be clear in section 2.4. Other major components, like the muon and hadron monitors, can be found in [8].

2.2.5 NuMI simulation

The geometry and physics associated with the NuMI beamline are implemented in a entirely GEANT4-based ([15]) Monte Carlo (MC) simulation called g4numi.⁷ The choice of GEANT4 is due its flexibility, allowing many user customizations, such us selecting between different physics models, attaching particular processes to particles and, even modifying the source code to extract desired information not usually available to users.

The origin of the simulation's coordinate system (called MCZERO) is located approximately at the front face of Horn 1 and along the trajectory of the primary proton beamline. The position convention follows right handed orientation Cartesian coordinates. The longitudinal axis (Z axis) direction overlaps with the beamline (i. e. 58 mrad downwards). The Y axis is vertical. Figure 2.8 shows the target in LE010 configuration (explained in section 2.3) and Horn 1 positioned in the MC coordinate system.

The simulation starts with a 120 GeV kinetic energy primary proton beam with a Gaussian beam transverse profile distribution ($\sigma = 1.1$ mm). The main geometrical details have been considered: the NuMI Target Hall including every component, the decay pipe, the absorbers, etc. However, the flux prediction is more sensitive to some elements than others. As described in section 2.5, any mismodeling, particularly

⁷The geant4 version used is geant4.2.p03.

concerning the focusing process (the target and horns) can lead to a bad prediction of the flux. The simulation chain ends at the point where a neutrino is produced.

we use the hadronic model package called FTFP_BERT, that combines the FRITIOF Precompound model [16] for processes with energies greater than 4 GeV and the Bertini cascade model for energies less than 5 GeV [17]. FTFP_BERT also incorporates the standard electromagnetic processes. More details about this model package and their agreement with measured data is discussed in chapter 4.

The g4numi output is a ROOT-based n-tuple file with a record of all information of the neutrino production. Neutrino generators like GENIE ([18]), used in MINERvA as the standard MC generator, use the meson decay points and momenta to generate neutrinos and simulate their interactions in a detailed detector geometry.

G4numi also calculates the probability of a neutrino to be produced in a particular direction given its parent kinematics. “Forcing” this direction changes the neutrino energy as well (as in equation 2.1). The probability of this occurring can be stored as a “weight” along with the new energy to make fast calculations of the flux in particular points in the beamline. The flux calculation at a single point is a good approximation for the flux at an extended detector if its location is far enough that sees the neutrino production as a point source. For detectors close enough to the neutrino production place to see it as an extended source, the flux at a single point in the detector is a rough estimation of the true flux through the detector, suitable for systematic uncertainty studies and to have a quick look of the neutrinos passing through a detector.

In this thesis, the flux for MINERvA (shown in chapter 5 to 7) is integrated over its fiducial volume (see section 3.5). The extension of our procedure (section 4) to other detectors takes NOvA Near Detector as example and the flux is calculated at the center of its front face.

Flux challenges

There are two main challenges when calculating NuMI flux using a simulation:

- **Hadron production.** The MC flux relies on the underlying hadronic models and their predictions can have significant disagreements with data since QCD is not a completely calculable theory at the energy scales of interest. Constraining hadron production models used in the simulation by applying external data is a major topic of this thesis and it will be cover extensively in chapter 4.
- **Focusing uncertainties.** Any geometrical mismodeling can lead to a bad flux prediction, especially around the target and horns system where the flux seems very sensitive to the hadron directions. The simulation has an idealized setup, where the implementation of the volumes needs often be simplified. On the other hand, the parameters involved in NuMI 's construction such as the location of beamline elements have inherently associated uncertainties. The MC simulation is used to calculate the effect of those uncertainties on the flux. This is covered in section 2.5.

2.3 NuMI fluxes

NuMI has been constructed as a tunable beamline. Different energy spectra and intensities of the neutrino flux can be achieved by changing the magnitude and direction of the magnetic field, as well as the relative distance among the target, Horn 1 and Horn 2. The idea is to be able to select and enhance the focusing of particles produced in a specific kinematic phase space and have a flexible beam energy. This is a key feature of NuMI that allows the exploration of neutrino oscillations at different Δm^2 (section 1.1).

The NuMI main modes are called Low Energy (LE) and Medium Energy (ME). During the LE era, the horns positions remain fixed while the target was remotely movable, allowing experimenters to tune the neutrino energy. As was mentioned in section 2.2.2, the target, Budal monitors and baffle were placed in the same carrier in order to make this change easily. For ME, the Horn 2 has been moved downstream from Horn1 while the target was moved upstream, producing the same energy shift in the flux spectrum but more efficiently as just moving the target.

The direction of the current pulsed to the horn conductors can be selected to focus or defocus hadrons with a particular charge. When π^+ (and K^+) are focused, it enhances the muon neutrino component of the flux. This mode is called “Forward Horn Current” (FHC). The opposite current direction creates a magnetic field that focuses mainly π^- to enhance the muon antineutrino flux. This mode is called “Reverse Horn Current” (RHC).

Low Energy Beams

As was mentioned, in LE the target is longitudinally movable and the current value can be selected. The convention for LE configurations is “LEXXXzYYYi”. The z=0 position (LE000z) corresponds the place where the target is inserted into

the Horn 1 as far as it can physically goes without touching the IC. XXX is the distance upstream from $z=0$ (in cm). The current is expressed as $YYYi$, where YYY is a reference magnitude (in kA) approximately the value used in the configuration.

Table 2.3 contains the LE runs during the MINERvA run period.

Name	Configuration	Current (kA)	Displacement from $z=0$ (cm)
LE FHC	LE010z185i	182.1	-10.0
LE RHC	LE010z-185i	-182.1	-10.0
LE OHC (“horn off”)	LE010z000i	0	-45.0
Pseudo ME FHC	LE100z200i	196.8	-100.0
Pseudo ME RHC	LE100z-200i	-196.8	-100.0
Pseudo HE FHC	LE250z200i	196.8	-250.0

TABLE 2.3: Naming convention and parameters for LE NuMI runs for MINERvA data period.

The LE010z185i and LE010z-185i configurations were the long term runs for the LE era (2005-2012) and they have been used to achieve the physics goals in MINERvA [4] and MINOS [19]. The main result of this thesis is the flux determination for these configurations. The target is positioned 10 cm from $z=0$ (as a visual aid, figure 2.8 a) shows the target location respect to the Horn 1 in the MC coordinate system, i.e, LE010z).

The black line in figure 2.12 shows the on-axis flux spectrum for ν_μ in “LE010z185i” with its characteristic peak around 3 GeV (“LE010z-185i” has similar peak for $\bar{\nu}_\mu$) and its long energy tail that is extended up to 120 GeV.

By moving the target upstream with respect to “LE010z” the peak energy shifts to higher values (table 2.3). The LE100z200i and LE100z-200i configurations have 100 cm target displacement from $z=0$ and their peaks are at 5 GeV for ν_μ and $\bar{\nu}_\mu$, respectively. In the “LE250z200i” configuration the target is moved 250 cm from $z=0$ and the peak shifts to 8 GeV. NuMI ran in these configurations for short periods of time allowing systematic cross-checks by measuring the neutrino interaction rates in MINOS and MINERvA in a given neutrino energy bin but with

different hadrons contributing in the different tunes. The spectrum when turning the horn currents off is also shown and we can note that the flux decreases significantly in this configuration.




FIG. 2.12: MC ν_μ fluxes for LE runs.

Figure 2.13 shows the yields of π^+ that leaves the target to generate neutrinos that pass through MINERvA in different NuMI configurations. For LE010z185i, most of the π^+ are focused in the the region between 2-15 GeV/c longitudinal momentum and transverse momentum less than 600 MeV. The biggest contribution comes from $p_Z = 7$ GeV/c as we expect for a 3 GeV LE peak using formula 2.1. For LE010z000i, just pions with very small transverse momentum will decay, rather than hitting material, to neutrinos that are forward enough to pass through MINERvA. This is the reason why the particle focusing is important in any modern conventional neutrino beam.

The LE100z200i and LE250z200i configurations, focus higher momenta π^+ for two reasons:




FIG. 2.13: π^+ yields that exit the target in for different beam configurations at MINERvA and NOvA Near Detector.

First, when the target is moved away from Horn 1, highly energetic pions that in LE pass neck-to-neck by the horns and decay to neutrino unlikely to travel on-axis can be focused by both horns now. This is the main contribution to the spectrum as can be seen in the “underfocused” plot in figure A.1 (appendix A).

Second, some “unfocused” and “horn2-only” pions are now part of the “horn1-only”, “underfocused” and “overfocused” categories. For example, a fast pion with small transverse momentum that is focused just by Horn 2 in LE can pass the Horn 1 with the same transverse momentum in the special runs (see figures A.2 to A.7).

Medium Energy Beam

By moving the target upstream, the peak energy of the on-axis neutrino flux increases. But this procedure is more efficient when the relative distance between horns is simultaneously enlarged. The reasoning is the same as in the “special runs”: more highly energetic neck-to-neck pions can be caught by the horns before their transverse momentum make them diverge from the beamline and crash into the walls. However, moving Horn 2 further away is more efficient than just moving the target. In this way, the ME flux spectrum is approximately 30% greater than in LE100z200i configuration.

The bottom left plot in figure 2.13 shows the π^+ yields off the target. A significant shift to larger longitudinal momentum values can be noted in comparison to LE010. The result is the flux spectrum in figure 2.14 (red line). The peak is approximately at 6 GeV with a small bump at 3 GeV due to overfocused pions.

The NOvA Near Detector is located 12.26 mrad off the beam axis and, due to equation 2.1, the neutrino energy spectrum is peaked at lower energy, and is significantly narrower than the spectrum seen on axis (see figure 2.14).





FIG. 2.14: MC ν_μ fluxes for ME runs in the MINERvA and NOvA Near Detectors. LE010185i is shown as a reference.


2.3.1 Beam components

Figure 2.15 shows the flux components for different beam configurations. The neutrino type focused in FHC modes are muon neutrinos with a small (few percent) background contamination from muon antineutrino, that is totally unfocused. The background comes mainly from mesons produced outside of the target and some “horn2-only” mesons. The electron neutrino flux in LE010z185i represents less than 1% of the total flux and about 0.8% in the focusing peak according to the simulation.

Conversely, muon antineutrinos focused in LE010z-185i flux are the main component up to 8 GeV. For higher energies (> 10 GeV), when the flux comes mostly from unfocused meson parents, the muon neutrino contribution is dominant due to the fact that more positive mesons are produced than negative mesons in the target but also K^+ are focused (see figure 2.15).



(a) LE010z185i and LE010z-185i beam configuration



(b) ME000z200i beam configuration

FIG. 2.15: Neutrino components for LE and ME beams.

2.4 Hadronic cascade in NuMI

When the primary proton beam interacts with the target (carbon) or with any other NuMI volume (made of carbon, aluminum, iron, helium, etc) it generates a hadronic cascade creating pion, kaons, protons and other particles. Some of them decay to make neutrinos but others may interact in any NuMI component. The hadronic interaction chain, i.e., the history of how a neutrino has been produced, is very important because each one of these interactions relies on a MC model and we need to know how well that model is representing nature.

The NuMI hadronic cascade is composed of many interactions of different particles and materials. The objective of this section is to recognize which of these interactions are relevant for the neutrino flux in order to look for data to apply in the correction procedure described in chapter 4. Most of the analysis is based primarily on ν_μ in LE010185i (with some ν_e plots). The corresponding plots for NOvA have been placed in appendix A.

2.4.1 Neutrino ancestries

Figures 2.16 a) and b) show the LE010z185i flux split in terms of the parent and grandparent identity for the main particle contributions that passes through MINERvA (plots NOvA are in figures A.9 and A.10). The parent identity plots are extended up 40 GeV but for simplicity, the corresponding grandparent plots are only shown to 10 GeV.

The main contribution for muon neutrinos at energies less than 20 GeV comes from π^+ ($\sim 97\%$). In terms of the grandparents, they come from the primary proton ($\sim 57\%$), secondary and tertiary protons ($\sim 15.8\%$) and π^+ ($\sim 10\%$). The high energy tail (ν energy > 20 GeV) is dominated by K^+ .

Electron neutrinos up to 10 GeV come, basically, from μ^+ decays ($\sim 85\%$) and

K^+ ($\sim 11\%$), and most of their grandparents are π^+ ($\sim 84\%$) For higher energies, the electron neutrino generation is dominated by K^+ with a small contribution of neutral kaons.

For NOvA, the trend is similar but the transition between parent identity is 6 GeV for ν_μ and 5 GeV for ν_e .


2.4.2 Interaction map

A tool has been developed to make a complete survey of all interactions which happened in NuMI that lead to a neutrino. This was done to obtain a more quantitative view of which projectile and produced particles, as well which materials, are relevant to understand the flux. This “interaction map” counts the average number of hadronic interactions per event for each neutrino flavor.


Figures 2.17 and 2.18 are examples of two interaction maps of ν_μ in LE010z185i integrated over 0-20 GeV (the corresponding NOvA maps are in figures A.11 and A.12). Fields with less than 0.001 interactions are not filled. The horizontal axes have the relevant NuMI materials.

The vertical axis of figure 2.17 has all possible hadron projectiles. The main contributions come from interactions that happen on carbon, basically in the target. Other relevant materials are aluminum (horn inner conductors), iron (decay pipe walls), and air (target hall). Other significant projectiles are π^+ and neutrons. Due to the neutrino energy cut at 20 GeV for these plots, the K^+ contribution does not appear.

Figure 2.18 shows the interaction map for incident proton. This map contains the number of produced particles per ν_μ events vs. the NuMI target materials (numbers less than 0.001 are not shown). Interactions with incident protons happen mainly in carbon, producing protons and π^+ .



(a) Parents.



(b) Grandparents.

FIG. 2.16: ν_μ and ν_e flux in the LE010z185i configuration at MINERvA split in their ancestry.




FIG. 2.17: Interaction map of incident particles vs. target material for ν_μ flux passing through MINERvA in the LE010z185i configuration (numbers less than 0.001 are not shown).




FIG. 2.18: Interaction map for proton projectiles. The map shows the number of produced particles per ν_μ event vs. target material at MINERvA in LE010z185i configuration (numbers less than 0.001 are not shown).

2.4.3 Hadronic production

The kinematic distribution of hadrons produced in primary proton interactions on the target is shown in figure 2.19 for 6 particles: π^+ , π^- , K^+ , K^- , protons and neutrons in terms of Feynman-x and transverse momentum. This corresponds to muon neutrinos in LE010z185i and similar plots for muon neutrino in NOvA are in figure A.13. For charged pions and kaons, an additional line on the top has been placed with the approximate expected neutrino energy using formula 2.1. The plots are normalized to 10^6 POT's and the bin sizes are: $\Delta x_F = 0.02$ and $\Delta p_T = 0.02 \text{ GeV}/c$.

The region of interest for π^+ is $x_F < 0.15$ and $0.05 < p_T < 0.55$. This region contributes to the focusing peak. π^- and K^- look defocused in the same region and their contribution is very small. K^+ production has an extensive area, but the region of interest is $x_F > 0.17$ where they become dominant in the neutrino spectrum. Protons have a wider x_F range and it can be split in two regions: one for $x_F < 0.95$ that has a peak around 0.5 and other for $x_F > 0.95$ that corresponds to protons likely to be quasi-elastics (see chapter 4). Neutrons make a very small contribution to the LE flux.

2.4.4 Beam attenuation

The relevant amount of material traversed by hadrons in the neutrino chain leading to a muon neutrino in LE010z185i is shown in figure 2.20 (Similar plot for NOvA can be found in figure A.14). The importance of this quantity in correcting the flux is seen in section 4.2.1. The figure corresponds to the primary and secondary protons passing through the target (C), and π^+ crossing the target (C), the horn inner conductors (Al), the decay pipe volume (He) and the decay pipe walls (Fe). The horizontal axis is the material traversed (mol/cm^2 units) and the bin size is




FIG. 2.19: Hadron production from the primary proton beam interacting in the target for ν_μ flux passing through MINERvA in the LE010z185i configuration.

0.1 mol/cm².

The top left plot corresponds to the primary proton in the target and shows the characteristic exponential behavior one expects. The peak at the 14 mol/cm² come from protons that leave the target without interacting. The top right plot shows the secondary protons per momentum in log scale with a peak at high momentum due to those created quasi-elastically.

The rest of the plots are for π^+ neutrino parents in different materials. In the target, most of them traverse less than 2.5 mol/cm² \sim 20 cm. Though the horns inner conductor is thin (2-4 mm), the horns are long and skinny resulting in rather long path lengths (around 1 mol/cm² \sim 10 cm). In the decay pipe it is 1 mol/cm² \sim 275 m but with a limit at 2.45 mol/cm² \sim 675 m, length of the pipe. The decay pipe walls are negligible.

Table 2.4 summarizes the density and mass number of the NuMI volumes included in figure 2.20. This corresponds to the volumes where the attenuation correction is applied (see chapter 4).

Volume	Material	Mass number (g/mol)	Density (g/cm ³)
TGT and BM	carbon	12.01	1.83
IC	aluminum	26.98	2.7
DVOL	Helium	4.003	0.000145
DPIP	iron	55.85	7.87

TABLE 2.4: Material information for volumes considered in figure 2.20.

2.5 Focusing uncertainties

In this section, I present the uncertainties in the flux that come from the focusing system. Some of these efforts were inherited from MINOS studies ([20]) like the transverse horn offset, the baffle scraping, the POT counting and the horn current magnitude. We revisited the horn current distribution and added the horn




FIG. 2.20: Material traversed of particles in the NuMI beamline for ν_μ flux passing through MINERvA in LE010z185i configuration.

inner conductor shape, target position and water layer around the inner conductor uncertainties.

We expect that those uncertainties directly related to the relation of target and horns as well as to the horn parameters impact the region just outside the focusing peak and has low impact for high energies. The reason for this is that all parameters involved (distances, materials, etc) have been optimized to have a small uncertainty on the flux peak (to be more precise, in the Far over Near ratio for oscillation studies) and there was big efforts to make the simulation very close to the actual setup in place. The biggest uncertainties are in the falling edge of the flux spectrum (4-6 GeV) as can be seen in the figure 2.21 that shows the fractional shifts by 1 sigma in the LE FHC mode.

A brief description of each uncertainty is presented in the rest of the chapter in the context of the ν_μ flux in LE010z185i at MINERvA and MINOS. These were calculated using MC simulation considering the uncertainty on each parameter given by the NuMI beamline group. The general procedure is to vary each parameter by $\pm 1, 2, 3\sigma$ around its accepted value and see the effect on the neutrino flux. Understanding an uncertainty means knowing, as well as we can, the effect of a change in that parameter. In our case, we need to see the effect on the flux when we varying a parameter. Some of the effects are clearly linear (like POT counting), while other effects need further investigation to see the exact functional dependency of the change. In this case, several sigma flux deviations were simulated and a second degree polynomial was used to find this form in different neutrino energy bins. The fractional shift on the flux is taken in each bin by evaluating the resulting polynomial at $\pm 1\sigma$.




FIG. 2.21: Focusing fractional shifts for ν_μ in the LE010z185i configuration at MINERvA.

2.5.1 Transverse horn offset

This uncertainty was considered to be a $\sigma = 1.0$ mm offset. In [20] Horn 1 and Horn 2 misalignments were study separately and the total effect is in the red line in figure 2.21. That line also include smaller effects as angular misalignments of the horns.

The result is the undesirable effect of reducing the flux in the focusing peak by 0.05% even while it increases by 2% the flux in the 4.5 - 8 GeV energy range. For example, in the focusing peak, “horn2-only” mesons with small angles that need slight focusing by Horn 2, are now deviated in Horn 1 and become unlikely to contribute to the flux when we move the Horn 2 (in analogy can be made for “horn2-only” mesons). But moving the horns makes high energy mesons that would pass neck-to-neck be focused and become more likely to contribute to the on-axis flux.

2.5.2 Baffle scraping

This uncertainty comes from neutrino ancestors that are produced when the primary proton beam interacts with the baffle upstream of the target First, by fitting the proton beam profile to a Gaussian and calculating the fraction of the beam expected to hit the baffle. Second, by measuring the beam induced temperature change in the baffle and relating that to the number of interactions. This gives a 0.25% protons that hits baffle as can be seen in [20].

Baffle scrapping decreases the number of low energy pions, including those that contribute to the focusing peak. On the other side, more high energy pions are able to be focused now. The orange line in figure 2.21 shows a very small reduction in 0-5 GeV neutrino energy and a flux increase of no more 1% in the 5-11 GeV region.

2.5.3 POT counting

NuMI directly measures the number of protons in the beam using two Toroid Intensity Monitor Integrators [8] with a uncertainty of 1% (see [20]). To take into account other beam related parameters like its position and width, the uncertainty is conservatively taken as 2%. The POT value is an overall normalization.

2.5.4 Horn current

The uncertainty on the horn current magnitude is estimated to be 1%, generating a corresponding uncertainty in the magnetic field value. The falling edge of the flux peak increases by 6.5% when the current increases (as seen in the green line in figure 2.21) and it decreases when the current decreases.

The skin depth effect (δ) of the conductors was reassessed [21] and it was found that the current is approximately uniformly distributed inside the conductor for $\delta \sim 7.7 \text{ mm}$ ⁸. We varied δ within its nominal uncertainty (due to the uncertainty on the magnetic permeability) and found no change in the flux, so the skin depth systematic was removed.

2.5.5 Horn inner conductor shape

The NuMI simulation was upgraded to incorporate a more accurate geometric and magnetic model of horn 1 [22].

The main addition is a better representation of the inner conductor shape. Due to some limitations in the geant4 geometry package, the parabolic part was approximated by segmenting it into in small cones with 50 μm length each one (it is called “old horn model” in this thesis). Ideally the horn model would have infinite

⁸This value correspond to 2.3 ms pulse duration and inner conductor thickness between 2-4 mm

number of infinitesimal length segments. The impact of improving the segmentation by making it 50 times finer (“new horn model”) and thus better modeled, is to reduce the flux up to 6% in the 4-6 GeV range for LE FHC and up to 14% in 7-14 GeV for ME FHC as can be seen in figure 2.22. This suggests that this change especially affects mesons leaving the target and passing the neck in their trajectory before decay.




FIG. 2.22: Effect of the new horn model on the ν_μ flux at MINERvA for the LE010z185i and ME000z200i configurations.

As it is described in section 3.5, the new horn model was not implemented for the current MINERvA published papers.

The new horn model has an effect that is very similar to the effect we see when we reduce the horn current by 0.8%. Using this fact (and due to the time scale of this thesis), we reused the horn current magnitude uncertainty shape for this

systematics assigning 100% uncertainty. The result is the light blue line in figure 2.21.

2.5.6 Longitudinal target offset

An optical survey has been made to determine the exact longitudinal position of the target after moving the target carrier between different NuMI configuration runs [23]. The uncertainties of these values are around 1 mm. Any shift from the nominal positions shown in table 2.3 affects the neutrino spectrum since the relative position target - Horn 1 is crucial in the particle focusing.

The precise position of the target was implemented in g4numi by splitting each NuMI beam configuration in run periods using the values given by the survey for the time when MINERvA starts taking data. Every run period is known as a playlist and it is named using the label **minervaX**, where X corresponds to a run period. This involves generating flux predictions for each playlist as can be seen in table 2.5.

Beam configuration	Playlist	ΔZ (cm)
LE010z185i	minerva1	+0.50
	minerva7	+0.82
	minerva9	-0.40
	minerva13	+0.87
LE010z-185i	minerva	+0.50
	minerva5	+1.15
	minerva10	+0.82
LE010z000i	minerva6	+0.82
LE100z200i	minerva2	+0.43
	minerva11	+0.83
LE100z-200i	minerva3	+0.43
	minerva12	+0.83
LE250iz200i	minerva4	+0.43
	minerva8	-0.09

TABLE 2.5: MINERvA LE flux playlists. The shift in the target position Δz is with respect to the nominal position.

Figure 2.23 shows the impact of the longitudinal target displacement in the ν_μ flux for LE FHC mode. Moving the target upstream produces a decrease in the flux in 4-6 GeV energy region.




FIG. 2.23: Longitudinal target position displacement effect in the ν_μ flux at MINERvA.

We use the longitudinal target position uncertainty from MINOS studies, that assume 1 cm uncertainty (blue line in 2.21) even when we use the survey data in our simulation. This value is clearly overestimated value and needs to be revised in the future.

2.5.7 Water layer uncertainty

The horns are constantly sprayed with cooling water which is deposited around the inner conductor creating a thin layer. The sketch in figure 2.24 shows the horn

inner conductor from the transverse view (left side) and the longitudinal view (right side) located between $a < r < b$ (light violet curve).




FIG. 2.24: Sketch of the position of the water layer around the horn inner conductor.

A 1 mm thick water layer was implemented in g4numi [24] (the NuMI Beam group estimates 1 mm thickness as the maximum expected) and the effect on the ν_μ flux in LE010z185i is shown in figure 2.25: four percent reduction on the flux to due the meson absorption on the water for energies less than 4 GeV. The uncertainty applied was 0.5 mm (blue line in figure 2.21).




FIG. 2.25: Effect of 1 mm water layer around the Horn 1 inner conductor in the ν_μ flux at MINERvA in the LE010z185i configuration.

CHAPTER 3

Overview of neutrino flux determination strategies

As was described in chapter 2, there are two primary sources of uncertainties in NuMI and in most conventional neutrino beams. One of them comes from the incomplete knowledge of the physics of the hadronic interactions that makes the simulation relies on phenomenological models, and the other one comes from some geometrical unknowns associated with the focusing process, relevant in some neutrino energy ranges. This chapter is dedicated to presenting different approaches to determine the neutrino flux in a conventional neutrino beam context as well the MINERvA strategy. We do not pretend to present the strategies of all experiment's approaches in detail. We want to place the MINERvA efforts as a part of a bigger effort in the experimental neutrino community.

We can distinguish two kind of procedures: those that determine the flux as an external input, i.e., without any intervention of the detector in where they are trying to find the flux (we show as an example the hadron production constraint using external data in section 3.1) and other procedures that require the use of

some data taken in the same detector for which we are trying to calculate the flux, but applying a criteria to deconvolute the flux (a “standard candle” process) that produced the data. This last procedure is called *in situ*, and we present three examples: multi-beam fitting (section 3.2), the low-nu method (section 3.3) and the neutrino - electron scattering constraint (section 3.4). Other procedures are not considered here (like the use of muon monitors [25]) because they are not within the scope of this thesis.

3.1 Using external data

To apply a constraint to the hadron production in the cascade that leads to a neutrino in the flux simulation, we need to have some knowledge about the number of hadrons produced in the interactions and their kinematic distributions.

During the past decades an extensive program has been running in many experiments to measure cross-sections of hadron-nucleus collisions in the non-perturbative QCD regime. This has been useful to understand the effects of the intra-nuclear interactions and test model hypotheses. In recent years, new experiments have been running, motivated by the search for the quark-gluon plasma and to help to improve the simulations of neutrino fluxes and cosmic ray air showers. Sections 4.3 to 4.7 contain lists of some of these experiments in the context of our hadron production correction procedure.

A complete and model independent knowledge of all interactions in this regime would mean having measurements of interactions of every particle with all incident energy possible on different targets. Even though there are many datasets available, there are still a number of areas that lack the data coverage needed for a specific conventional neutrino beam. Recent experiments dedicated to measuring the production of hadrons from proton beams incident on thin (few percentage of

interaction length) Be, C, Al targets. These experiments were optimized to measure hadrons which make a large contribution to the neutrino flux in running experiments like MiniBooNE, T2Km and MINOS. Other efforts involved to make direct measurements on the same target (or in a replica) used to produce a neutrino beam. The idea is to use all data available, incorporating their uncertainties and making well-founded arguments to extrapolate to areas not covered by data.

The MiniBooNE experiment used data from the HARP experiment at CERN to determine their flux. HARP measured the cross-sections of proton and pions with momenta range from 1.5 GeV/c to 15 GeV/c interacting in different target materials like beryllium, carbon, aluminum, etc. ([26], [27], [28], [29], and [30]) and also measured the yields of particle production in a MiniBooNE replica target ([31]) with a good forward coverage.

The Booster beamline (see section 2.2.1) collides 8 GeV protons into a 71.1 cm long Be target ([32]), and, in conjunction with a one magnetic horn system, creates an on-axis muon neutrino beam peaked at 500 MeV - 600 MeV. Figure 3.1 a) shows the kinematic distribution of π^+ at the target that contributes to the ν_μ flux at the MiniBooNE detector. The figure was taken from MiniBooNE flux prediction paper [33]. The black box represents the HARP coverage of π^+ produced from proton - Beryllium interactions at 8.9 GeV [29] that was used to constrain the flux. Most of the pions relevant to MiniBooNE are constraint by the HARP data. More details about other datasets incorporated by MiniBooNE and their procedure can be found in their flux paper mentioned before [33].

Another example is the T2K flux constraint using NA61 data. Their muon neutrino beam is peaked at 0.6 GeV and is produced by colliding protons at 30 GeV with a 91.4 cm long graphite target [34]. NA61 measured the cross-sections of protons interacting at 30 GeV in carbon nuclei ([35] and [36]) and a T2K replica target ([37]). Figure 3.1 b) shows the π^+ kinematic distribution at the target that con-

tributes to the muon neutrino flux at the T2K far detector (Super-Kamiokande). The area enclosed by the black lines corresponds to the NA61 π^+ data coverage that was used in the flux calculation ([34]) showing that most of pion production is able to be constrained. A complete description of their procedure and all datasets used can be found in their flux paper [34].


3.2 Using a multi-beam fitting

As was explained in section 2.3, changing the beamline configurations may change the neutrino spectra. This allows us to relate, in each configuration in the simulation, the neutrino yield in one energy bin and a kinematic region of the hadrons produced in the target. Using the data measured in these different configurations, we can attempt to find the hadron yields from the target that make data and simulation agree for all configurations considered. This is an example of a “*in situ*” procedure.


MINOS followed this strategy ¹ using data from different beam configurations at their Near Detector, adjusting parametrized yields of the hadrons that exit the target [20]. The functional form used was similar to the BMPT parametrization [38]. Their goal was to find the best extrapolation of the flux at the Near Detector to the flux at the Far Detector [1] for their oscillation parameter search. In the introductory chapter, we used figure 1.2 to highlight the importance of the flux determination. That figure shows how the F/N ratio changes due to the beam fit procedure.

The result of MINOS’s beam fit procedure is shown in figure 3.2. The beam configurations have the same convention as we explained in section 2.3 where the data and simulation before tuning and the simulation after the tuning are repre-

¹In addition to the hadron production, MINOS also adjusted some beam focusing parameters.



(a) π^+ neutrino parents in MiniBooNE [33]. The black box enclosed the data coverage of HARP data of π^+ produced in proton - beryllium interactions at 8 GeV ([29]).



(b) π^+ neutrino parents in T2K [34]. The area enclosed by black lines corresponds to the data coverage of NA61 π^+ produced in proton - carbon interactions at 30 GeV ([35]).

FIG. 3.1: Examples of hadron production experiments for neutrino flux (HARP and NA61).

sented by dots, a gray line and a black line respectively. The effects on the event rates MINOS Near Detector can be seen in the ratios.




FIG. 3.2: Results of the multi-beam fitting procedure followed by MINOS to correct the NuMI flux in different configurations [39].

3.3 Low-nu method

The differential cross-section of the neutrino-nucleon charged current scattering respect to the recoil energy (ν) can be expressed as a combination of integrals over structure functions (A, B and C) and the ratio ν/E_ν , where E_ν is the neutrino energy, as shown in equation 3.1:

$$\frac{d\sigma}{d\nu} = A \left(1 + \frac{B}{A} \frac{\nu}{E} - \frac{C}{A} \frac{\nu^2}{E^2} \right) \quad (3.1)$$

The cross-section becomes independent of the neutrino energy as the ratio between the recoil energy and the neutrino energy goes to zero ($\nu/E_\nu \rightarrow 0$) (see [40] for more details). The energy spectrum of events with small ν/E_ν will approximate the energy dependence of the flux (the flux shape). An additional normalization to external is required (for instance, a cross-section measurements), the flux can be determined.

The method was developed by the CCFR/NuTeV collaboration to determine the neutrino flux at energies greater than 30 GeV and it used this flux to calculate the neutrino cross-sections [41]. on iron. Some efforts have been made to extend the procedure to lower energies by MINOS and MINERvA. MINOS used this procedure to calculate the muon neutrino and antineutrino inclusive cross-section on iron ([42] and [39]).

Recently, MINERvA also estimated the flux in the LE010z185i configuration using the low-nu method [43]. There are two challenges associated with the estimate and the procedure used to make it:

- To find a correction for a finite ν values. MINERvA uses GENIE to compute corrections and the uncertainty on them.
- To accurately reconstruct the hadronic recoil energy.

The low-nu MINERvA results are used in this thesis and additional details are shown in section 6.

3.4 Neutrino - electron scattering constraint

Another *in situ* procedure uses elastic neutrino scattering on electrons as a standard candle. Standard electroweak theory predicts this cross-section precisely since only leptons are involved in the $\nu e \rightarrow \nu e$ process, as shown in figure 3.3.




FIG. 3.3: neutrino - electron scattering Feynman diagram.

The $\nu e \rightarrow \nu e$ scattering process produces a single forward electron and a neutrino. The electron can be easily identified with a high detector resolution like MINERvA. However, the cross-section is approximately 2000 times smaller than compared with the neutrino - nucleon cross-section and is thus statistically limited. The neutrino produced escapes the detector and it is not possible to reconstruct the incident neutrino' energy. However, it serves as a flux normalization since the number of interactions can be counted and we can then the number of neutrinos that passed through the detector.

Recently, MINERvA studied these events [44] and this is applied as an additional constraint for the flux determined in this thesis. More details are provided in

section 6.

3.5 Minerva strategy

MINERvA has adopted a multi-layer approach to predict the NuMI flux that involves all the procedure described before: the use of external hadron production data (section 3.1), the low-nu method (section 3.3), the neutrino-electron constraint (section 3.4) and a multi-beam fitting of the hadron production(section 3.2).

The MINERvA flux prediction has been constantly updated throughout, motivated by several factors like the incorporation of new hadron production data, a better understanding of the the effect of flux systematics, and a geometrical improvements in the NuMI simulation. The first two flux versions, called Generation 0 and Generation 1, have been used as the standard flux in the current MINERvA publications (see later table 5.1). The most updated flux version - Generation 2 - will be used in all upcoming MINERvA papers and it is the basis of the final results in this thesis. The details about every generation will be provided in the next chapters.

The MINERvA strategy is the following:

- The foundation is formed from constraining hadron production with external measurements on thick and thin targets (see chapter 4). The use of independent data gives us the chance to have different predictions.
- Beam optics uncertainties are incorporated by propagating errors in the alignment of beamline elements and the horn's geometry and magnetic field (see section 2.5). The flux determined by hadron production constraints plus the focusing uncertainties is called *a priori* flux.
- In-situ measurements are then incorporated atop the foundation. These consist

of:

- The rate of charged current muon neutrino and antineutrino scattering events with low energy transfer to the target for which the flux shape is known (see section 3.4). It is used to study the consistency of our flux predictions.
- Neutrino-electron scattering events for which the cross-section is known (see section 3.4). It is used to reduce the flux uncertainty.
- The Multi-beam fitting technique had been followed early on before MIPP NuMI data was published and it is not used in this thesis. However, it served to check sources of MINERvA reconstruction issues.

One criteria used to design the MINERvA strategy has been the desire to develop flux calculating tools that can be shared with other NuMI experiments. That is the reason for **PPFX** (see section 4.8).

The next chapter explains the hadron production correction procedure we followed and the incorporation of the datasets we used in the context of Generation 2 flux version ² (see section 4.3 to 4.7).

The details of Generation 0 and Generation 1 are postponed to chapter 5 when a historical review is made and the *a priori* LE flux results of every generation is presented. Chapter 6 is dedicated to the incorporation of *in situ* measurements and to compare Generation 0-1 to Generation 2 in the way to update the MINERvA results already published. In chapter 7, we will extend our procedure to ME flux and other detectors.

²Explaining the dataset incorporation in the context of Generation 2 seems more convenient since it is the base of this final thesis results

CHAPTER 4

Correcting the simulation using hadron production data

The non-perturbative QCD nature of the hadronic interactions in any conventional neutrino beam requires the use of phenomenological models in the simulation. In principle, if we have enough experimental data to constrain the model, the flux can be determined and its uncertainty can be well understood. As was explained in the introductory chapter, knowing the flux is important in neutrino oscillation studies and it is crucial for neutrino cross-section measurements. How can the NuMI beam flux simulation be constrained using hadron production data? This chapter addresses this question by developing the MINERvA strategy introduced in section 3.5.

This chapter is structured in three main parts. The first one establishes some common definitions (section 4.1) and explains in detail the procedure we followed to correct the hadron production (section 4.2). In the second part we survey all existing relevant data for NuMI and describe their implementation in the context of Generation 2 flux version (sections 4.3 to 4.7). The last part is dedicated to intro-

ducing the tool we developed to handle the data into a computational framework: PPFX (section 4.8).

4.1 Definitions

Before explaining our procedure, it is worthwhile to check some basic definitions that we use in this chapter and throughout the rest of the thesis.

4.1.1 Total cross-section

In hadron - hadron interactions ($\mathbf{h+h}$), the total hadronic cross-section (σ_{total}) can be divided in two components, one elastic ($\sigma_{elastic}$) and the other one, inelastic ($\sigma_{inelastic}$):

- **Elastic component** ($h + A \rightarrow h + A$): the nucleus remains intact after the interaction. This implies that neither mesons nor nucleons are ejected from the nucleus.
- **Inelastic component** ($h + A \rightarrow + \text{mesons} + \text{fragments of nucleus}$): the nucleus is broken and new particles are created, like mesons.

However, when we consider hadron - nucleus interactions ($\mathbf{h+A}$) another component arises: “quasi-elastic”:

- **Quasi-elastic component** ($h + A \rightarrow + p + \text{fragments of nucleus}$): the nucleus is broken but no new particles (mesons) are created. It is characterized by a fast nucleon in the final state (typically, $x_F > 0.95$) and one or more slow nucleus fragments.

In the context of $\mathbf{h + A}$ interactions, we define the **absorption** cross-section ($\sigma_{absorption}$) as the sum of the quasi-elastic and inelastic components. Then, the total

hadronic cross-section is:

$$\sigma_{total} = \sigma_{elastic} + \underbrace{\sigma_{inelastic} + \sigma_{quasi-elastic}}_{\sigma_{absorption}} \quad (4.1)$$

4.1.2 Differential cross-section

The **Lorentz invariant differential cross-section** is defined as:

$$f = E_s \frac{d^3\sigma}{dp^3}. \quad (4.2)$$

where E_s and p are the energy and momentum of the particle produced in the interaction.

The approximate scaling variable “Feynman-x” ([45] and [46]) is defined as:

$$x_F \equiv \frac{p_{\parallel}^*}{P_{\parallel}^*(max)} \simeq \frac{2p_{\parallel}^*}{\sqrt{s}} \quad (4.3)$$

where $(*)$ indicates that the variable is calculated in the center of momentum system, p_{\parallel}^* is the longitudinal momentum of the produced particle and $P_{\parallel}^*(max) \simeq \sqrt{s}/2$ is the maximum momentum allowed (\sqrt{s} is the energy of the center of mass). Feynman speculated that the cross-sections of inclusive high energy hadronic collisions an energy independent scaling behavior when expressed in terms of x_F .

4.2 Procedure

This section is dedicated to explaining in detail the procedure we followed to make the hadron production correction.

Each interaction in the neutrino’s ancestry chain is corrected with a weighting factor computed from yields or invariant differential cross-sections:

$$c_i = \frac{N_i^{data}}{N_i^{MC}} \quad (4.4)$$

The factor f_i depends on the identities of the projectile, target, and produced hadron as well as the initial and final state kinematics. Here “i” denotes the initial and final state information, and it is generally a particular bin in a hadron production dataset. In that case we have an uncertainty σ_i and a covariance with other bins j, V_{ij} . Even in the case that the interaction in question is not covered by a hadron production dataset, we still assign c_i , σ_i , and V_{ij} factors using our best judgment. N_i^{MC} is either taken directly from the MC or computed by simulating the interactions in question to produce a yields table or an invariant cross-section.

We refer to each f_i as the “central value” estimate for the interaction. In section 4.2.5 we will describe the way in which the V_{ij} are used to vary the weighting factors to propagate uncertainties.

In addition to the weights applied to each interaction we also account for attenuation of particles in the beamline, inserting correction factors which depend on the absorption cross-section in data and MC.

4.2.1 Attenuation Corrections

When a beam particle passes through a volume, the interaction rate depends on the cross-section and the amount of material traversed. The probability $P(r)$ of a particle not interacting while crossing a distance r is given by:

$$P(r) = e^{-rN_A\rho\sigma} \quad (4.5)$$

Where N_A is the Avogadro number, ρ is the nuclei volume density and σ is the absorption cross section per nucleus.




FIG. 4.1: Particle beam traversing a volume. Left side: particle leaving a volume without interacting. Right side: particle interacting in volume.

If an interaction does not happens (see figure 4.1, left side), this correction is applied to the probability that the particle survived:

$$c(r) = e^{-r \frac{N_A \rho (\sigma^{data} - \sigma^{MC})}{A}} \quad (4.6)$$

The material traversed $r N_A \rho$ is independent of the specific material and it can be expressed as mol/cm². The correction is shown in the figure 4.2 considering a cross-section disagreement up to ± 20 mb and up to 10 mol/cm² of material.




FIG. 4.2: The survival correction.

There is no correction to apply when the agreement between data and MC is perfect (the factor in this case is 1). When the material traversed is very small the

factor is also close to 1, for example, when the particle is created on the edge of a volume. If the MC overestimates the cross-section, less particle beam survives in the simulation and the correction goes up and increases for higher material traversed. Conversely, if the MC underestimates the cross-section, more particle beam survived in the simulation and the correction goes down and even decreases for higher material traversed.

If an interaction happens in a volume (see figure 4.1, right side), the correction is applied to the probability that the particle survived up to the interaction point:

$$c(r) = \frac{\sigma^{data}}{\sigma^{MC}} e^{-r \frac{N_A \rho (\sigma^{data} - \sigma^{MC})}{A}} \quad (4.7)$$

Now, the attenuation correction also depends on the data over MC cross-sections ratio. The figures 4.3 and 4.4 show this correction for two fixed MC cross-section values: $\sigma = 193.6$ mb (this corresponds to π on C at 7 GeV) and $\sigma = 368$ mb (π on Al at 7 GeV) respectively. As before, in a perfect data-MC cross-section agreement the correction would be 1. Interactions are more (less) likely to occur in the simulation if the MC overestimates (underestimates) the cross-section and this effect competes with the survival part of the function creating the patterns we see in the figures.

4.2.2 Particle Production

Thin target experiments tend to release their data as invariant double differential cross-sections f in terms of Feynman-x x_F , transverse momentum p_T . As was mentioned in section 4.1.2 Feynman-x is an approximate scaling variable and it lets us to extrapolate a cross-section taken at one energy to another one in inelastic collisions. When scaling data like NA49 that was taken at 158 GeV (see section 4.5.1) we use FLUKA ([47], [48]) to remove the residual energy dependence and scale to




FIG. 4.3: Attenuation correction when an interaction happened for $\sigma_{MC} = 193.6 \text{ mb}$ (this corresponds to π on C at 7 GeV).




FIG. 4.4: Attenuation correction when an interaction happened for $\sigma = 368 \text{ mb}$ (this corresponds to π on Al at 7 GeV).

12-120 GeV ([49]) that is the range relevant for NuMI. This “energy scaling” correction is shown in figure 4.5 for four x_F ranges for positive pions generated in proton on carbon collisions. For typical p_T values (~ 300 MeV/c) the correction at 120 GeV is very small (less than 1%) but it can be up to 10% when the energy of the incident proton goes down to 12 GeV.




FIG. 4.5: FLUKA energy scaling correction for invariant cross-section data taken at 158 GeV to lower energy down to 31 GeV. This is split in four x_F ranges.

The formula to be applied becomes:

$$c(x_F, p_T, E) = \frac{f_{Data}(x_F, p_T, 158\text{GeV}) \times scale(x_F, p_T, E)}{f_{MC}(x_F, p_T, E)} \quad (4.8)$$

The energy scaling correction has been calculated for the following incident energies: $\{12, 20, 31, 40, 60, 80, 100, 110, 120\}$ GeV and the MC cross sections f_{MC} for $\{12, 20, 31, 40, 50, 60, 70, 80, 90, 100, 110, 120\}$ GeV. A simple linear

interpolation is applied for intermediate incident energies. In figure 4.6 the blue line represents NA49 data integrated in p_T and expressed in terms of the angle of the pion produced as in NA61. The green line shows the effect of applying the energy scaling factors and shows that it is in good agreement with NA61 (purple line), especially around the focusing peak as it is shown in a zoom version at the right plots.

This prescription was checked by scaling NA49 pion production data at 158 GeV to NA61 data taken at 31 GeV. The difference between the two was very small. The effect on the the neutrino flux was calculated to be no more than 1% (see [50]). The figures in 4.7 show the fractional difference between NA49 and NA61, where $\Delta f(x_F, p_T)$ is the discrepancy between the value of scaled NA49 value at 31 GeV and the corresponding NA61 value.

When an interaction happens in a volume where we are applying an attenuation correction, we have to modify equation 4.8 to cancel the inelastic cross-section contribution in the formula 4.3 to avoid a double correction:

$$c(x_F, p_T, E) = \frac{f_{Data}(x_F, p_T, 158\text{GeV}) \times scale(x_F, p_T, E)}{f_{MC}(x_F, p_T, E)} \times \frac{\sigma_{MC}(E)}{\sigma_{data}(E)} \quad (4.9)$$

On the other side, thick target experiments release their data as yields of particle production when they leave the target ¹. The correction in this case is given by the ratio of the data over MC yields:

$$correction(p_Z, p_T) = \frac{n_{Data}(p_Z, p_T)}{n_{MC}(p_Z, p_T)} \quad (4.10)$$

¹Longitudinal dependence of the yields is sometimes provided like in [37].





FIG. 4.6: Effect of the energy scaling for π^+ produced on proton - carbon collisions. The corresponding x_F and p_T are indicated in the left side plot and the right plot is a zoomed version.





FIG. 4.7: Fractional difference ($\Delta f(x_F, p_T)/f(x_F, p_T)$) between NA49 and NA61 due to energy scaling (taken from [50]).

Details of the target attenuation correction when using thick target data

Applying both the thick target correction and the target attenuation correction gives a longitudinal target dependence to the hadron production correction. This is important since the hadrons that exit the target are focused differently depending on how they enter the horns. However, the challenge, as in the case of thin target correction, is to avoid double counting. The solution is not simple because there are many particles interacting or surviving in the target.

As we are going to see in section 4.7, among all particles passing through the target, only proton attenuation correction has the central value different than 1 and particularly, the primary proton beam has the most significant contribution in comparison to any secondary proton produced in the target. Pions and other particles do not change the central value.

Having this in mind, we assume that when a hadron that leaves the target is able to be corrected with thick target data, it is a good approximation to apply an attenuation correction to just the primary beam and consider that all other particle attenuation corrections are embedded in the thick target measurements.

Another simplification is that we consider only the longitudinal material traversed to calculate the attenuation correction since the primary proton momentum is high and almost parallel to the beamline, and the target length is also parallel to the beamline longitudinal axis.

However, two modifications are needed to make this possible. The first one is to calculate $n(p_Z, p_T)$ per interaction instead of per proton incident on the target by using a factor y , that is the fraction of the primary proton beam that interacts in the target. In this way, the beam that passes through the target without interacting can be treated independently. Equation 4.10 becomes:

$$\boxed{c(p_Z, p_T) = \frac{n_{Data}(p_Z, p_T)/y_{Data}}{n_{MC}(p_Z, p_T)/y_{MC}}} \quad (4.11)$$

The second modification preserves the thick target data yield we are applying. The MC yield for a (p_Z, p_T) bin can be expressed in terms of the longitudinal distribution $g(z)$ of the primary proton interaction positions in the target as:

$$n_{MC} = \int_0^{z_{max}} f(z) dz \quad (4.12)$$

where z_{max} is the maximum longitudinal distance traveled by the proton in the target before interacting. After applying the attenuation correction c_{att} , the n_{MC} becomes:

$$n'_{MC} = \int_0^{z_{max}} g(z) c_{att}(z) dz \quad (4.13)$$

And then,

$$n_{MC} = n'_{MC} \times \frac{n_{MC}}{n'_{MC}} \quad (4.14)$$

$$= \int_0^{z_{max}} g(z) [c_{att}(z) \times \frac{n_{MC}}{n'_{MC}}] dz \quad (4.15)$$

The factor n_{MC}/n'_{MC} is calculated per bin and the final form of the equation 4.7 is:

$$\boxed{c(r) = \frac{\sigma^{data}}{\sigma^{MC}} e^{-r \frac{N_A \rho (\sigma^{data} - \sigma^{MC})}{A}} \times \frac{n_{MC}}{n'_{MC}}} \quad (4.16)$$

4.2.3 Extending data coverage

Based on theoretical guidance, some extension of data is possible to different materials and different incident and produced particles.

The first extension is “material scaling” and we are looking here for an uncertainty associated with using hadron production data taken on carbon in other materials. As was exhibited by Barton et al. [51], the invariant cross-section can be parametrized as:

$$f(A, x_F, p_T) = \sigma_0 A^{\alpha(x_F, p_T)} \quad (4.17)$$

where A is the mass number of the target nuclei different than hydrogen². The parametrization is independent of the energy of the incident particle as the approximate scaling x_F hypothesis suggests. Using this approach, the relation between the cross-sections of two materials can be express as:

$$\frac{f(A_1, x_F, p_T)}{f(A_2, x_F, p_T)} = \left(\frac{A_1}{A_2} \right)^{\alpha(x_F, p_T)} \quad (4.18)$$

MINERvA has followed the parametrization of α proposed by [51] and extended by [38]:

$$\alpha(x_F, p_T) = (a.x_F^2 + b.x_F + c).(d.p_T^2 + e.p_T + 1) \quad (4.19)$$

$\alpha(x_F, p_T)$ is found by doing an independent fit of the Skubic data [52] of invariant cross-sections of K^0 , Λ^0 and $\bar{\Lambda}^0$ produced from proton interacting in various nuclei at 300 GeV. Figure 4.8 shows an example of the $\alpha(x_F, p_T)$ fit for neutral kaon production in proton interactions at $x_F = 105 \text{ GeV}/c$ and angle = 0.9 mrad. The same fit is made with all Skubic data, having 318 α values in total.

The results are checked against π^\pm and K^\pm production in proton interactions to find the additional uncertainty to cover materials different than carbon with proton carbon data (more details an be found in [53]).

²The data fit using the equation 4.17 in [51] excludes hydrogen.




FIG. 4.8: Skubic data $\alpha(x_F, p_T)$ fit for neutral kaon production in proton interactions at $x_F = 105$ GeV/c and angle = 0.9 mrad. The same fit is made with all Skubic data. This is shown as example (taken from [53]).

The additional uncertainties to extend data taken in carbon to other materials (p_T units are GeV/c) are in table 4.1.

Another extension comes from the isoscalar nature of ^{12}C (The main material of the NuMI target). The isospin symmetry of deuterons establishes that $\sigma(pd \rightarrow \pi^+ nd) = \sigma(nd \rightarrow \pi^- pd)$ and then we can infer that:

$$\sigma(pC \rightarrow \pi^\pm X) = \sigma(nC \rightarrow \pi^\mp X) \quad (4.20)$$

This symmetry is used to determine the correction for hadron production in neutron carbon interactions and assume that the same correction is necessary for other materials than carbon. No additional uncertainty is added when using this extension. Since there is a small number of neutron interactions in the typical neutrino ancestry chain, and because we could not find any relevant data to test equation 4.20.

(a) π^+ .				(b) π^- .			
$x_F \backslash p_T$	[0, 0.5]	[0.5, 1]	[1, 1.5]	$x_F \backslash p_T$	[0, 0.5]	[0.5, 1]	[1, 1.5]
[0, 0.25]	10	7.5	2.5	[0, 0.25]	12.5	7.5	2.5
[0.25, 0.5]	10	12.5	10	[0.25, 0.5]	10	7.5	15
[0.5, 0.75]	7.5	10	2.5	[0.5, 0.75]	15	10	20

(c) K^+ .				(d) K^- .			
$x_F \backslash p_T$	[0, 0.5]	[0.5, 1]	[1, 1.5]	$x_F \backslash p_T$	[0, 0.5]	[0.5, 1]	[1, 1.5]
[0, 0.25]	10	7.5	2.5	[0, 0.25]	12.5	7.5	2.5
[0.25, 0.5]	10	12.5	10	[0.25, 0.5]	10	7.5	15
[0.5, 0.75]	7.5	10	2.5	[0.5, 0.75]	15	10	20

TABLE 4.1: Additional uncertainties (%) when extending data taken in carbon to other materials(p_T units are GeV/c).

The neutral kaons can be determined from charged kaons assuming isospin symmetry in a quark parton model [38]. If the number of valence quarks u is the double of d and $q_s = \bar{q}_s$ for $q = \{u, d, s\}$, then:

$$N(K_{L(s)}^0) = \frac{N(K^+) + 3N(K^-)}{4} \quad (4.21)$$

In this way a correction for neutral kaons for any material can be calculated when K^\pm data is available and we also extend also this procedure to thin target data. Since the kaon data is dominated by statistical errors, no uncertainty is added from these assumptions.

4.2.4 Procedure when there is no data

We have two options in assigning an uncertainty to the hadron production when there is not clear theoretical guidance to extend the data coverage.

1. First we can examine the spread between different hadronic models. This was the original approach in MINERvA (see sections 5.1 and 5.2) to handle interactions

not covered by NA49. The differences between models could depend on the type of interactions. Then, to avoid underestimating the true flux differences predicted by each model, we categorized interactions not covered by NA49 and looked at the maximum difference between geant4 hadronic models for the following categories each individual interaction that is relevant for the total neutrino flux and assumes this as the uncertainty. A complete list of categories and more details can be found in [54]. Examples of these categories are:

- π^+ neutrino parents that were born in the target whose ancestry includes a proton reinteractions in the target.
- K^+ neutrino parents that were born inside the magnetic horns.

Figure 4.9 shows the predicted fluxes due to π^+ neutrino parents born in the target whose ancestry involves a previous pion interaction from 5 geant4 hadronic models. The spread in this case is $\sim 40\%$ around the average value in the focusing peak.




FIG. 4.9: Example of geant4 model spread: neutrino flux from π^+ parents born in the target with a pion ancestry.

The problem with this procedure is that it is hard to determine any internal

correlation between models and we end up with the risk of underestimating the final flux error.

2. Second we apply a best guess uncertainty guided by the agreement with other datasets. Our idea is that we are not totally blinded for interactions that there is not data. The MC hadronic model we used (FTFP_BERT) is a micro-physical model that applied QCD (in general, particle physics) first principles to all components in the interactions. Then we expect that the agreement (or disagreement) between data and MC for interactions that have data should be roughly the same as interactions not covered by data.

We notice three characteristics of the corrections when we look at the data:

- the corrections between different produced particles are not correlated.
- most of the correction values are in 0.6-1.4.
- the corrections depend on x_F in a first order approximation, i.e., not p_T dependence, since the neutrino energy is proportional to x_F .

Considering this, for the interactions where we do not find any data or extension to apply, we categorized those interactions in 4 x_F regions (0-0.25, 0.25-0.5, 0.5-0.75 and 0.75-1.0) per any combination of projectile and produced particle and assign 40% uncertainty and treat them uncorrelated each other. This procedure will be applied in section 4.6.1. Some examples of these categories are:

- $\pi^+ A \rightarrow \pi^- X$ with x_F in 0.25-0.5 .
- $\pi^+ A \rightarrow \pi^- X$ with x_F in 0.5-0.75 .
- $pA \rightarrow pX$ with x_F in 0.75-1.0 .

4.2.5 Uncertainty propagation

We use the so called “multi-universe” technique to propagate uncertainties ([55]). This technique is the creation of a statistical ensemble of individual randomly generated universes, where each “universe” selects a value (deviate) for each uncertain parameter. The parameters for the neutrino flux determination are the hadron production cross sections, yields, etc. The deviates are drawn from a multi-variable normal distribution centered on the parameter’s central values:

$$N(\vec{x}|\vec{\mu}\mathbf{V}) = \frac{1}{(2\pi)^{M/2}\det(\mathbf{V})^{1/2}} \exp(-\frac{1}{2}(\vec{x} - \vec{\mu}) \cdot \mathbf{V}^{-1} \cdot (\vec{x} - \vec{\mu})) \quad (4.22)$$

where \mathbf{V} is the covariance matrix of all M parameters, $\vec{\mu}$ is the vector of the central values and \vec{x} is the vector of deviates. \mathbf{V} is decomposed by the Cholesky factorization in a lower triangle \mathbf{L} and an upper triangle matrices:

$$\mathbf{V} = \mathbf{L}\mathbf{L}^T \quad (4.23)$$

Then, the vector of deviates \mathbf{x} is given by ³:

$$\vec{x} = \vec{\mu} + \vec{R}\mathbf{L} \quad (4.24)$$

where \vec{R} is vector of random Gaussian distribution values with mean zero and unit variance. Every universe predicts a flux spectrum and the resulting spread from the statistical variations is used to evaluate the final flux systematic uncertainty .

Figure 4.10 shows an application of the multi-universe technique for the flux using 50 universes applying just the MIPP NuMI pion production weights and assuming +75% bin-to-bin correlation for the systematic errors (more on this in section 4.4.1).

³The proof of this procedure can be found in the Section 7.4 of ([56])




FIG. 4.10: Example of hadron production uncertainty propagation using the multi-universe technique.

The final correction for a particular universe to the neutrino flux is applied as the product of all corrections given to the neutrino ancestry information.

4.3 Data available

We developed our procedure for correcting the simulated flux in the previous section. We now look into the hadronic cascade in NuMI (section 2.4), to make a survey and find all relevant data that can be used in the NuMI beamline.

Thick target data

These are experiments that collide monochromatic protons on a thick target and measure the yields of hadrons leaving the target. This the first place to look since they are designed to help the flux determination efforts by using a similar primary beam and the same or replica target of the neutrino beamlines. In our

survey, we found two datasets:

- MIPP: proton on a spare NuMI target at 120 GeV:
 - π^\pm for p_Z up to 80 GeV/c ([57]).
 - K/π for $p_Z > 20$ GeV/c ([58]).
- NA61: π^\pm production from proton collisions on a replica T2K target at 31 GeV ([37])⁴.

Thin Target Data

These are experiments that use monochromatic beams on targets of a few percentage interaction lengths. They measure:

- The inelastic and absorption cross section. Some of the datasets found are:
 - Belletini et. al. ([59]), Denisov et al. ([60]), etc. : proton, pions and kaons on carbon, aluminum, etc. in a wide energy range.
 - NA49 ([61]): proton on carbon at 158 GeV.
 - NA61 ([35]): proton on carbon at 31 GeV.
- Hadron Production. Some of the datasets found are:
 - Barton et. al ([51]): $pC \rightarrow \pi^\pm X$ at 100 GeV for $x_F > 0.3$.
 - NA49 ([61]): $pC \rightarrow \pi^\pm X$ at 158 GeV for $x_F < 0.5$.
 - NA49 ([62]): $pC \rightarrow n(p)X$ at 158 GeV for $x_F < 0.95$.
 - NA49 ([63]): $pC \rightarrow K^\pm X$ at 158 GeV for $x_F < 0.2$.
 - NA61 ([35]): $pC \rightarrow \pi^\pm X$ at 31 GeV.

⁴This data is not used in this thesis but it can be used as a cross check in the future

- MIPP ([64]): π/K from pC at 120 GeV for 20 GeV/c.

In the next four sections, we explain some details of the experiments and their use in the hadron production correction for the NuMI flux determination, focused on those parts that are relevant to understand their results and uncertainties, as well as the assumptions of our implementation in the computational framework (section 4.8). Thin target data is split in three sections, one dedicated to the hadron production data (section 4.5), the second one dedicated to the data extension (section 4.6) and the other one dedicated to the absorption and inelastic cross-section data (section 4.7).

4.4 The thick target datasets

The thick target data we used come from the experiment MIPP (**M**ain **I**njector **P**article **P**roduction) that is dedicated to studying cross-sections and yields of hadron production using thin and thick targets with multiple particle beams at Fermilab. One of its goals was to measure yields of hadrons that exit the NuMI target with low uncertainties in order to help determine the NuMI flux for Fermilab neutrino experiments. 1.43 million events were collected colliding 120 GeV protons from the Main Injector beam on a spare NuMI target. A schematic view of MIPP detector can be seen in figure 4.11 (more details can be found in [57] and [58]).

The basic setup consists of several sub-detectors, each one of which provides particle identification (PID) for a specific momenta range of the particles when they leave the target:

- a Time Projection Chamber (TPC) covering 0.2-1.2 GeV/c.
- a Time of Flight (ToF) covering 0.5-2.5 GeV/c.
- a segmented gas Cherenkov detector (Ckov) covering 2-20 GeV/c.

MIPP

Main Injector Particle Production Experiment (FNAL-E907)




FIG. 4.11: Schematic view of MIPP detector [57].

- a gas ring imaging Cherenkov (RICH) covering 4-80 GeV/c.

For the NuMI target run analysis, the particle identification relies on the highly segmented TPC and RICH detectors that give good momentum coverage. The ToF was used to estimate backgrounds in the TPC. The Ckov was not used in this analysis⁵. The low momentum ($p_Z < 2 \text{ GeV}/c$) PID is handled by the TPC while high momentum PID ($p_Z > 20 \text{ GeV}/c$) is handled by the RICH.

To highlight the importance of these data, we quantify the interactions covered by using thick target measurements. Figures 4.12 and 4.13 show the average number of inelastic interactions per π^+ and per K^+ per MIPP bin, starting from the primary proton beam interaction and counting up all inelastic interactions up to the creation of the hadron that exits the target. They are separated into three momentum ranges of the particles exited the target: 0-2 GeV/c, 2-20 GeV/c and 20-80 GeV/c. The

⁵It is complicated for ToF and Ckov to distinguish between particles when there is high multiplicity events.

hashed bins represent regions where there are no data due to detector acceptance. For K^+ , only the high momentum is relevant due to the data coverage (see section 2.4). The minimum possible value is 1 since meson production requires having the generation of new particles from proton-nucleus collisions. Numbers larger than 1 indicate reinteractions in the target. Many of these are incident pions and kaons that we have no thin target data to use in a correction (see next section).

There are more than 2 interactions on average for very low momentum pions (< 1 GeV/c) and these values decrease monotonically for higher momentum particles approaching 1.2 in the upper limit of MIPP coverage. Figure 4.12 shows that a significant number of interactions are not covered while the high energy pions in figure 4.13 are well covered.


4.4.1 Thick target pion production

MIPP NuMI results of π^+ and π^- yields ([57]) are presented in 124 and 119 bins respectively for p_Z in 0.3-80 GeV/c and p_T in 0-2 GeV/c per proton on target (POT). Figure 4.19 shows data published for π^+ (left side) and π^- (right side). Different markers represents different p_T ranges. Each p_T range has been multiplied by a factor to accommodate all the data points in a single plot.


Figures in 4.15 show a combination of the statistical and the background systematic uncertainties for π^+ and π^- . Most of the values are small and typically in 2-5% except for the edges of the sub-detectors, where the errors become bigger. In our approach, we treat these uncertainties just as statistical, considering them to be uncorrelated between bins.

The additional MIPP systematic uncertainties are:

- **Momentum scale** when a correction is applied to the reconstructed momenta to account for energy loss, scattering and any reconstruction algorithm bias. The



(a)



(b)

FIG. 4.12: The importance of secondary interactions: these plots show the average number of inelastic interactions per π^+ that exits the target using MIPP binning for p_Z in 0-20 GeV/c according to our NuMI simulation. p_Z and p_T are the momentum of the π^+ when it exited the target. The hashed bins represent regions where there are no data.





FIG. 4.13: The importance of secondary interactions: these plots show the average number of inelastic interactions per π^+ and K^+ that exit the target using MIPP binning for p_z in 20-80 GeV/c according to our flux simulation. p_z and p_T are the momentum of the π^+ and K^+ when they exited the target. The hashed bins represent regions where there are no data.




FIG. 4.14: MIPP charged pion production measured at NuMI (taken from [57]). The left side for π^+ and the right side for π^- . Different markers represents different p_T range. Every yield has been multiplied by a factor to accommodate them in single plots.

uncertainty is 1% in most of the bins.

- **Bin migration** due to the momentum resolution and reconstruction failures (4%) and mismodeling of the noise (1%).
- **Detector Modeling** in the MC (2%).
- **Pileup** due to an improper modeling of overlapping tracks, specially for high multiplicity events. This is the dominant uncertainty, typically a few percent but it can be up to 10% for some bins.

The figures in 4.16 show these additional systematics errors added in quadrature for π^+ and π^- . Nearly all bins are in 4-5% but large bin to bin correlations within each sub-detector are expected.

The effects of using four different positive bin to bin correlation assumptions of their systematic errors are shown in figure 4.17. The fractional errors on the flux





FIG. 4.15: MIPP NuMI charged pion statistical uncertainties. Numbers in every bins indicate the exact uncertainty value. The hashed bins represent regions where there are no data.





FIG. 4.16: MIPP NuMI charged pion systematic uncertainties. Numbers in every bins indicate the exact uncertainty value. The hashed bins represent regions where there are no data.

correspond to the part of the ν_μ flux that is affected by MIPP (i.e., the neutrinos that have one pion correctable by MIPP). As we expect, when we increase the correlation, the fractional error increases accordingly. For example, the values go from $\sim 2\%$ for no correlation scenario to $\sim 5\%$ for a totally correlated case around 8 GeV neutrino energy.




FIG. 4.17: Effect of the bin-to-bin correlation of the systematic uncertainties. The fractional error comes from systematic uncertainties applied to the ν_μ flux affected by MIPP assuming 4 possible correlations: none, +50%, +75% and +99%.

Technical details of the implementation

We MIPP NuMI charged pion correction uses the equation 4.11. The implementation is as follows :


1. As was mentioned above, we treat the statistical + background systematic uncertainties as just statistical. In that way, they enter as uncorrelated to the additional systematics presented.

2. A positive 75% correlation is used between bins for the systematic uncertainties within TPC and RICH. Bins from different sub-detectors are uncorrelated.
3. We have not made any data interpolation and the correction is applied directly when the kinematics of a pion falls into the bin edges.
4. To calculate the factor y_{Data} in equation 4.11, we look at the percentage of primary protons that pass through and leave the target without interacting in MIPP NuMI is $13.5 \pm 2 - 3\%$. This is based on the number of reconstructed tracks per proton incident on the target and assumes that one track events are just primary protons ⁶. This uncertainty comes from the reconstruction efficiency. We use 3% uncertainty for the uncertainty on this value.
5. In the MC, we found that 13.28% of primary protons leave the target without interacting or after a quasi-elastic interaction and then we use this value to calculate y_{MC} .


Figures in 4.18 show the MIPP charged pion weights using the formula 4.11. The Z-axes are restricted to $[0.5, 2.0]$ since all weights are inside this range. For low momentum, the MC underestimates most of π^+ yields for TPC bins. For RICH, there are two clear regions, one of them where MC overestimates the yields ($10 \text{ GeV}/c < p_Z < 30 \text{ GeV}/c$ and $p_T < 0.4 \text{ GeV}/c$) up to 25%, and another one where the π^+ are overestimated for $\sim 50\%$.

For π^- weights, the TPC bins have similar patterns as π^+ , however, for RICH, they are dissimilar. Apart for $20 \text{ GeV}/c < p_Z < 50 \text{ GeV}/c$ and $p_T < 0.4 \text{ GeV}/c$ where the MC overestimates π^- , the rest of the bins have small corrections. These will cause different effects in the ν_μ than $\bar{\nu}_\mu$ fluxes.

⁶These values were establish after an email exchange with J. Paley (main author of [57]).



(a) π^+ .



(b) π^- .

FIG. 4.18: MIPP NuMI charged pions weights.

4.4.2 Thick target kaon production

We extract kaon production by combining pion yields from [57] and the K/π ratios reported in [58] for longitudinal momentum ($p_Z > 20 \text{ GeV}/c$). For the ratio analysis, only the RICH was used and the results are presented in 24 bins.




FIG. 4.19: MIPP π^-/π^+ , K^-/K^+ , K^+/π^+ and K^-/π^- ratios taken from [58]. The momentum bin convention (X,Y) is such that X represent the longitudinal momentum and Y the transverse momentum. X(GeV/c) = 0 (20-24), 1 (24-31), 2 (31-42), 3 (42-60), 4 (60-90). Y(GeV/c) = 0 (<0.2), 1 (0.2-0.4), 2 (0.4-0.6), 3 (0.6-1.0), 4 (>1.0).

Figure 4.20 shows the statistical errors of K^+/π^+ and K^-/π^- and as can be noted, the values are high, especially in the latter case.

The systematic uncertainties on π/K were evaluated independently from the pion yield analysis presented above and we do not have enough information to cancel





FIG. 4.20: MIPP NuMI statistical uncertainties for K/π . Numbers in every bins indicate the exact uncertainty value. The hashed bins represent regions where there are no data.

any correlation between them. The main sources on the systematic uncertainties on the ratio are:

- **Beam:** estimating the effect of small changes of the proton beam profile on the ratio ($<4\%$).
- **Momentum:** accounting for the differences between the measured momentum in data with respect to MC. The effect on the ratio is $\pm 5\%$ for most of the bins.
- **Background subtraction,** typically less than 20%.

Figures in 4.21 show the systematic errors added in quadrature for K^+/π^+ and K^-/π^- . The values are big in comparison to pions, especially for high longitudinal momentum.

Technical details of the implementation

The implementation of the MIPP NuMI kaon correction is as follows:

1. The K/π statistical and systematic uncertainties are considered as uncorrelated. No bin to bin correlation assumption is made for the systematic uncertainties on the ratio.
2. No interpolation is applied to the data.
3. The K^\pm yields are determined by multiplying the π^\pm yields and K^\pm/π^\pm ratios in each universe.
4. $K_{S(L)}^0$ correction is calculated by using the equation 4.21.

Figures in 4.22 and 4.23 show the MIPP charged and neutral kaon weights using formula 4.11 after the kaon yield determination. The Z-axes are restricted to $[0.5, 3.0]$ but K^+ weights can be higher for $p_Z > 56 \text{ GeV}/c$. Except for $p_Z <$







FIG. 4.21: MIPP NuMI statistical systematic for K/π . Numbers in every bins indicate the exact uncertainty value. The hashed bins represent regions where there are no data.

40 GeV/c and $p_T < 0.4$ GeV/c the MC underestimates the K^+ yields. K^- and neutral weights have, in general, smaller values than K^+ .




(a) K^+ .




(b) K^- .

FIG. 4.22: MIPP NuMI charged kaons weights.



(a) K_L^0 .



(b) K_S^0 .

FIG. 4.23: MIPP NuMI neutral kaons weights.

4.5 The thin target hadron production datasets

The thin target hadron production datasets we used come essentially from NA49. Barton and MIPP are used in conjunction with NA49 to extend its data coverage.

4.5.1 NA49

The NA49 experiment's goal is to make a comprehensive study of soft hadronic interactions using different beams and thin targets. Their relevant results for NuMI are the measurements of hadron production in proton-carbon interactions at 158 GeV/c beam momentum. 377,000 inelastic events in the H2 beam line at CERN SPS accelerator complex were recorded from a beam composed by 65% protons, 30% pions and 5% kaons. Charged pions and protons invariant differential cross section as well as neutron yields were published in a wide x_F coverage([61] and [62]). There is also low x_F charged kaon data from a MINOS thesis [63].

A schematic view of the NA49 detector can be seen in figure 4.24 (more details can be found in [65]):

The basic setup consists of four large Time Projection Chambers (TPC):

- **Two Vertex TPC's** just downstream of the target, together with two aperture Vertex Magnets, are used for interaction vertex, particle tracking and PID determination.
- **Two Main TPC's** are located downstream of the magnets in order to extend the acceptance to higher momentum and provide sufficient track length for PID.

The NA49 interaction trigger system consisted of a small scintillation counter (2 cm in diameter) located 380 cm from the target and between the vertex TPC's (figure 4.24). This operated in anti-coincidence with the primary beam.




FIG. 4.24: Schematic view of NA49 detector [66].

The main systematics in NA49 come from:

- **Normalization** in determining the inelastic cross-sections from the interaction trigger. A trigger scintillator counter was placed in anti-coincidence with the beam. This systematic accounts for any losses of protons, pions and kaons when they hit the trigger as well as subtracting the elastic component. The uncertainty is 2.5% and it is directly applicable to the differential cross-sections. For neutron yield, it is 1.5%. Normalization is the largest contribution to the total systematics and it is expected to be fully correlated bin to bin.
- **Feed-down** induced by weak decays of strange particles, especially important for pions and protons (1-2.5%) as well as neutrons (3%). This is the second largest contribution to the systematics. For kaons, it can be neglected. It is expected to be uncorrelated bin to bin.
- **Binning** is the correction for finite bin width determining the deviation of the real cross-section at the center of the bin from the measured one. It is approximately 0.5%.

- **Trigger bias** due to fact that secondary produced particles can hit the trigger and cause the event to be vetoed. This is approximately 0.5-1%.
- **Tracking efficiency:** 0.5%.
- **Other uncertainties** like detector absorptions, pion decays and reinteractions in the target are in total about 0.5%.
- **Other neutron uncertainties** related to the reconstruction.

A summary of the NA49 systematics is shown in the table 4.2:

particle produced	total systematic (quadratic sum)	total systematic (upper limit)
π^\pm	3.8%	7.5%
K^\pm	4.2%	8.7%
backward protons	4.7%	10.0%
forward protons	3.7%	7.0%
neutrons	10%	28.0%

TABLE 4.2: NA49 systematics uncertainties.

4.5.2 Barton et al.

The result reported by [51] (Barton et al.) is a comprehensive cross-section measurements of fast secondaries (momentum of the particles produced in 30-88 GeV/c) with different incident beams (pions, kaons and protons) at 100 GeV using the Fermilab Single Arm Spectrometer facility in the M6E beamline (see page 741-775 in [67] for a description). The use of the same detector to study the A dependence of the invariant differential cross-sections cancels some of detector performance systematic uncertainties.

The relevant Barton data for NuMI is the pion production from proton - carbon interactions at 100 GeV. However, there is a disagreement of the pion and proton production between Barton and NA49 in the data points where they overlap. We

are going to explain later in this section the assumptions we made to use Barton data.

4.5.3 MIPP thin target

Ratios of high energy K/π have been studied in [64] from proton - carbon interactions. As will be explained later, combining NA49 and MIPP thin target data allows us to extend the data to higher energy kaon production.

4.5.4 Thin target pion production

NA49 data is used to correct the pion invariant differential cross-sections in the $-0.1 < x_F < 0.5$ and $0 < p_T < 2$ GeV/c range. Figure 4.25 shows some of the 270 bins per particle of NA49 data taken from [61] (the right plot corresponds to π^+ and the left one to π^-). Many data points populate the x_F region below 0.15 that are not shown in the plots. The lines come from NA49 interpolation (we implement our own interpolation as described later).

Barton et al. ([51]) reported their pion invariant differential cross-section data for x_F in 0.3-0.88 and $p_T = 0.3, 0.5$ GeV/c. As was pointed out, Barton and NA49 are in disagreement in points where they overlap. Figure 4.26 (upper plot) shows both datasets for $0.3 < x_F < 0.5$ (MC values in the plot are just for reference). Barton cross-sections for π^+ are approximately 25% less than NA49 as discussed in [61] (similar differences are seen for π^-). However, the ratios π^+/π^- agree between both datasets. We use this fact to extend the NA49 data for $x_F > 0.5$ by imposing a 25% reduction to Barton data and adding a 25% uncertainty due to this assumption. Figure 4.26 (lower plot) also show π^+ Barton data for $x_F > 0.5$.






FIG. 4.25: NA49 charged pion production in proton carbon interactions.



(a) Barton - NA49 comparison for π^+ in $0.3 < x_F < 0.5$.



(b) π^+ Barton data for $x_F > 0.5$.


FIG. 4.26: Barton et al and NA49 π^+ data in proton - carbon interactions.

Technical details of the implementation


The implementation of the pion production correction is as follows:

1. The NA49 and Barton data are interpolated to smooth transitions between bins. The granularity is $\Delta x_F = 0.005$ and $\Delta p_T = 0.025 \text{ GeV}/c$. This is finer than the data binning. MC values have the same granularity.
2. The statistical and systematic uncertainties assigned to each bin is the closest fractional error value of NA49 or Barton datasets.
3. The statistical uncertainties are assumed to be uncorrelated. Figures in 4.27 show these uncertainties for the interpolated NA49 data. The region for low x_F (0-0.25) and $p_T < 0.8 \text{ GeV}/c$ has small errors (less than 5%) for both π^+ and π^- . This is important to determine the flux focusing peak of MINERvA and NOvA as can be seen in chapters 5-7.
4. The systematic uncertainties are assumed to be 100% correlated between bins. It is clearly an overestimated value since only the normalization uncertainty is known to be fully correlated. A computational complication leads to this decision: the calculation of the Cholesky decomposition of a big matrix (for instance, in the case of NA49: 121 x_F bins \times 81 p_T bins makes a 9801-dimension matrix to be handled ⁷. Equation 4.23 demands that we compute the Cholesky decomposition of the covariance matrix. The total systematic uncertainties (see table 4.2) added in quadrature is used.
5. An energy scaling correction is applied to use these datasets for any incident proton on carbon in $12 - 120 \text{ GeV}/c$.

⁷Some techniques can be used to deal with this complication but a simple assumption had to be made due to the timescale to present the results of this analysis.



(a) π^+ .



(b) π^- .

FIG. 4.27: NA49 statistical error for pion production.

Figures in 4.28 show the weights for π^+ and π^- applying equation 4.8. For $x_F < 0.3$, π^+ below $p_T = 0.4 \text{ GeV/c}$ are underestimated by the MC and overestimated otherwise. For π^- , almost the entire region for $p_T < 1 \text{ GeV/c}$ has to correct the MC up.

Pion production from neutron - carbon interactions was also corrected with the NA49 and Barton data using equation 4.20.

4.5.5 Thin target kaon production


NA49 data is used to correct the invariant differential cross-sections for x_F in $[0, 0.2]$ as reported by [63] (figures in 4.29 a) and b) for K^+ and K^- , respectively). The lines are interpolated values in [63] (we make our own interpolation). Except for low p_T , the coverage is good for low x_F kaons.

A combination of NA49 pion ([61]) production with MIPP K/π ([64]) is used to extend the kaon data beyond $x_F = 0.2$. Figure 4.30 shows the MIPP kaon ratios (we are interested in the right hand side plots). The binning convention is described in the figure caption.


Technical details of the implementation

The charged kaon production correction uses the equation 4.11 as for charged pions case and its implementation is as follows:

1. As for pion production, NA49 kaon production data is interpolated to smooth transitions between bins. The granularity is finer than the data binning: $\Delta x_F = 0.005$ and $\Delta p_T = 0.025 \text{ GeV/c}$. However, no interpolation is made for MIPP pC ratios. MC values have the same granularity as their corresponding data after interpolation.




(a) π^+ .




(b) π^- .

FIG. 4.28: Thin target weights for charged pions.



(a) K^+ .



(b) K^- .

FIG. 4.29: NA49 invariant differential cross-section of charged kaon production in proton carbon interactions (taken from [63]).





FIG. 4.30: MIPP π^-/π^+ , K^-/K^+ , K^+/π^+ and K^-/π^- ratios taken from [64]. The momentum bin convention (X,Y) is such that X represent the longitudinal momentum and Y the transverse momentum. X(GeV/c) = 0 (20-24), 1 (24-31), 2 (31-42), 3 (42-60), 4 (60-90). Y(GeV/c) = 0 (<0.2), 1 (0.2-0.4), 2 (0.4-0.6), 3 (0.6-1.0), 4 (>1.0).

2. The statistical and systematic uncertainties assigned to each NA49 bin is the closest fractional data error. For MIPP, the data uncertainty is used directly.
3. The statistical uncertainties are assumed to be uncorrelated. The NA49 statistical uncertainties for K^+ and K^- production are shown in figures in 4.31. The values are typically higher than pions, as we expect. The statistical uncertainties for MIPP K/π ratios are in figures at 4.32 with the exact value written in each bin.
4. In consistency with pion correction, the NA49 systematic uncertainties are assumed to be 100% correlated between bins. The total of the systematic uncertainties (see table 4.2) added in quadrature is used for NA49. The figures in 4.33 show the systematic uncertainties used in MIPP thin target K/π ratios. We do not assume any bin to bin correlation for MIPP data ratios.
5. The energy scaling correction is applied for any incident proton on carbon in $12 - 120 \text{ GeV}/c$.
6. Formula 4.21 is used to estimate the neutral kaon cross-section from charged kaon data. The MC K_L^0 and K_S^0 yields are calculated off-line.
7. The factors y_{Data} and y_{MC} are calculated as for charged pion case.


Figure 4.34 a) and b) and shows that the weights for K^+ and K^- respectively. Neutral kaons weights are shown in figures at 4.35).

4.5.6 Thin target nucleon production

Proton production in pC interactions has been measured in NA49 [62] as invariant double differential cross sections for $-0.8 < x_F < 0.95$ and $p_T < 2 \text{ GeV}/c$. Protons produced quasi-elastically have Feynman-X greater than 0.95 and they are




(a) K^+ .




(b) K^- .

FIG. 4.31: NA49 statistical uncertainties for charged kaon production.




(a) K^+/π^+ .




(b) K^-/π^- .

FIG. 4.32: MIPP pC statistical uncertainties for K/π .




(a) K^+/π^+




(b) K^-/π^-

FIG. 4.33: MIPP pC systematic uncertainties for K/π




(a) K^+ .




(b) K^- .

FIG. 4.34: Thin target weights for charged kaons



(a) K_L^0 .



(b) K_S^0 .

FIG. 4.35: Thin target weights for neutral kaons.

not part of this data since those events are always vetoed by NA49 trigger counter. Figures in 4.36 show the data for proton (left side) and anti-proton production (right side) taken from [62]. Just the proton cross-section correction has been used since the anti-protons have very small effect. The values for negative x_F are not shown in the figure and the lines correspond to the NA49 interpolation reported in the paper. We have made our own interpolation.

NA49 also measured neutron production yields (dn/dx_F) integrated over the p_T acceptance for $0.1 < x_F < 0.9$. Figure 4.37 shows the 8 data points released by NA49 ([62]).

Technical details of the implementation

The implementation of the nucleon production correction is as follows:

- Proton production data has been interpolated with granularity $\Delta x_F = 0.005$ and $\Delta p_T = 0.05 \text{ GeV}/c$. The neutron yield was interpolated with a 4th degree polynomial and sliced in bins with granularity $\Delta x_F = 0.005$.
- As before, the statistical and systematic uncertainties assigned to each NA49 bin after interpolation are the closest fractional data uncertainty.
- Figure 4.38 a) shows the statistical uncertainties of proton production. The values are typically less than 5% for $p_T < 1 \text{ GeV}/c$. For neutron yield, the statistical error is around 10-15% (figure 4.38 b).
- The total systematic uncertainties added in quadrature (table 4.2) and 100% correlation bin to bin are assumed. For protons, it is necessary because the high number of bins that enter into the Cholesky decomposition. For neutrons it is not needed but it is made anyway for consistency with the other datasets




FIG. 4.36: NA49 invariant differential cross-section for proton production in proton carbon interactions (taken from [62]).




FIG. 4.37: NA49 neutron production yields in proton carbon interactions (taken from [62]).


used. Figures in 4.38 a) and b) show the fractional uncertainties for protons and neutron production, respectively.

- As in pion and kaon, energy scaling correction is used for nucleon production to go from 158 GeV incident proton incident to $12 - 120 \text{ GeV}/c$.


The central part of the proton production cross-section correction is overestimated and has to be corrected down as can be seen in figure 4.39a. The neutron yield needs to be corrected down for $0.15 < x_F < 0.65$ and up otherwise figure 4.39b.

4.6 Thin target data extension in Generation 2

This section deals with the cases when we do not have data to apply directly in Generation 2. As was described in section 4.2.3, the first approach can be to look



(a) proton



(b) neutron

FIG. 4.38: NA49 nucleon statistical uncertainties





FIG. 4.39: Thin target weights for nucleon production.

if any theoretical guided data extensions and then use our best guess anchored in data (see section 4.2.4). In any case, Generation 2 tries to avoid the use of hadronic model spread as a procedure for these cases.

4.6.1 Interactions of nucleons not covered by thin target data

For interactions with nuclei other than carbon, we can apply the thin target carbon data from the previous section following the procedure described in section 4.2.3. The material scaling adds the uncertainty listed in table 4.1 to the data uncertainty for charged pions and kaons with $0 < x_F < 0.75$ and $p_T < 1.5 \text{ GeV}/c$. Another extension we used is to apply proton interaction data to the neutron interactions using the isospin symmetry principle (section 4.2.3).

For any other nucleon interaction not covered yet, we use our best guess uncertainty based on data as was described in section 4.2.4, i.e, 40% uncertainty split in four x_F uncorrelated regions in 0-1 (0-0.25, 0.25-0.50, 0.50-0.75 and 0.75-1.0) per nucleon incident: {proton, neutron} and per particle produced: $\{\pi^+, \pi^-, K^+, K^-, K^0, \text{proton, neutron, any other particle}\}$. This gives 64 different uncertain parameters, treated as uncorrelated. One thing that should be noted is that the quasi-elastic interaction correction falls into this category.

4.6.2 Incident mesons and other interactions

Currently, we are not applying any data corrections for interactions with incident mesons. In these cases, we use 40% uncertainty for each 4 uncorrelated x_F regions (0-0.25, 0.25-0.50, 0.50-0.75 and 0.75-1.0). We do this for each different combination of incident meson: $\{\pi^+, \pi^-, K^+, K^-, K^0, \text{any other meson}\}$ and produced particle: $\{\pi^+, \pi^-, K^+, K^-, K^0, \text{proton, neutron, any other particle}\}$. This gives

144 possible uncorrelated uncertain parameters.

When an interaction is not covered by any data and any of the above categories, a correlated 40% uncertainty is applied. Examples of these interactions are $pC \rightarrow \bar{p}X$. As we see in the next chapters, the contribution from these “others” kind interactions is very small.

4.7 Thin target inelastic and absorption cross section datasets

4.7.1 Proton - carbon inelastic cross-section

Figure 4.40 shows the data and MC inelastic cross-sections comparison for proton and neutron carbon interactions. The datasets are Bellettini et al [59], NA61 for T2K [35], Denisov et al. [60], Carroll et al. [68], NA49 [61] and Roberts et al. [69]. Some inconsistency in terminology in Denisov et al leads us to assume that the data provided in the paper is the absorption cross-section instead of the inelastic cross-section. For that reason, we subtract the quasi-elastic part using the Roberts et al calculation based Glauber model. The neutron cross-section is placed as reference. The data suggests that the inelastic cross-section has no energy dependence up to 300 GeV incident energy.

The MC inelastic cross-sections are shown with black markers. They have been calculated by running high statistic thin target simulations of proton incident with different energies and looking at interactions where no new particles (mesons) are created. The MC has a nonphysical behavior, increasing its value monotonically when the energy is increased. However, the MC absorption cross-section is flat at 243.24 mb.

4.7.2 π^\pm absorption cross-sections

Figure 4.41 shows the absorption cross-section for charged pion incident in carbon and aluminum up to 70 GeV/c momentum. The datasets used are Cronin et al [70], Denisov et al [60], Allaby et al [71], Allardyce et al [72], Longo et al [73], Vlasov et al [74], Bobchenko et al [75], Carroll et al [68]. The data suggest that the absorption cross-sections decrease monotonically after 1 GeV/c incident momentum.




FIG. 4.40: Proton-carbon inelastic cross sections.

The MC values are in blue (pion on carbon) and red (pion on aluminum).

The data - MC discrepancies can be seen in figures 4.42 a) and b). The differences between data and MC absorption cross-sections are displayed. The data has a small disagreement with MC. For pion on carbon, most of the differences are less than 10 mb and for pion on aluminum, less than 20 mb, except at low energies.

4.7.3 K^\pm absorption cross-sections

The figures in 4.43 show the absorption cross-section for charged kaon incident in carbon and aluminum up to 60 GeV/c momentum. The datasets used are Abrams et al. [76], Denisov et al [60], Carroll et al [68] and Allaby et al. [71]. There are few data points for K^+ and no pattern has been found (see figure 4.43 a)). For K^- , a slow cross-section decrease in energy can be seen in figure 4.43 b).

Figures 4.44 and 4.45 show the difference between data and MC absorption cross-sections for K^+ and K^- on carbon and aluminum. The disagreement is higher




FIG. 4.41: π C and π Al data and MC absorption cross-section.

in comparison to charged pions, especially at low energy.

4.7.4 Technical implementation of the absorption and inelastic cross section

The correction is implemented as follows:

1. pC inelastic and absorption cross sections:

- For the inelastic cross-sections (needed in cases when we use equation 4.9), we implement the value given by NA49 data with its associated uncertainty ($226 \text{ mb} \pm 2.5\%$). For the absorption cross-section (needed for the proton attenuation correction in the target), we add the implemented inelastic cross-section plus the quasi-elastic component that has 40% uncertainty as described in section 4.6.1.





FIG. 4.42: π Data - MC absorption cross-section data - MC comparisons. Error bars on many of the points are too small to be seen.





FIG. 4.43: Charged kaon on C and Al data and MC absorption cross-section.





FIG. 4.44: K^+ data - MC absorption cross-section data - MC comparisons.





FIG. 4.45: K^- data - MC absorption cross-section data - MC comparisons.

- The MC, the calculated inelastic cross-section for the incident proton energy shown in figure 4.40 is implemented and making a linear interpolation for intermediate incident proton energies.
- We should note that the this correction has a central value different than 1.

2. π^\pm and K^\pm absorption cross section:

- The central value for this correction is equal 1.
- For π^\pm , we noted that almost all differences between the datasets and MC are less than $\pm 5\%$ of the MC value. Based on this, the uncertainty associated to $\pi^\pm\text{C}$ and $\pi^\pm\text{Al}$ are 10 mb and 23.8 mb, respectively.
- For K^\pm , the data and MC discrepancies are higher than π^\pm , specially for low momentum ($< 2\text{ GeV}/c$) and this discrepancies are different for K^+ and K^- . Based on this, we implement uncertainties that cover most of the tries almost all dataset - MC in figures 4.44 and 4.45. Table 4.3 lists these values.

3. Details about attenuation correction:

- This correction is applied to all hadron neutrino parents, grandparents and when it exist, great-grandparents.
- For the target, the amount of carbon traversed longitudinally by the particles interacting or leaving the target is tabulated precisely (we add the material traverse only in the fins).
- The correction is also applied to others NuMI components: the horn inner conductor (Al), the decay volume (He) and decay pipe walls (Fe).
- For nucleons and in general any other particles, we assign the biggest uncertainty (K^+ on aluminum).

A summary of the uncertainty per particle and material is listed in table 4.3:

data	$\Delta\sigma$ (mb)
π on carbon	10
π on aluminum	23.8
K^+ on carbon ($P < 2 \text{ GeV}/c$)	68
K^- on carbon ($P < 2 \text{ GeV}/c$)	80
K^+ on aluminum ($P < 2 \text{ GeV}/c$)	83
K^- on aluminum ($P < 2 \text{ GeV}/c$)	49
K^+ on carbon ($P > 2 \text{ GeV}/c$)	13
K^- on carbon ($P > 2 \text{ GeV}/c$)	20
K^\pm on aluminum ($P > 2 \text{ GeV}/c$)	30.5

TABLE 4.3: Summary of the meson absorption cross-section uncertainty implemented in out hadron production procedure.

4.8 PPFX

Note: This is a technical section which may not interest some readers.

PPFX is the package we wrote to implement the hadron production corrections and propagate uncertainties described in the previous section. PPFX stands for **P**ackage to **P**redict the **F**lu**X** and is an experiment independent neutrino flux determination package for the NuMI beam that provides a correction for hadron production mismodeling⁸ using almost all relevant data. Currently, PPFX corrects the beamline simulated with g4numi using geant4.2.p03 and the FTFP_BERT hadronic model⁹. The inputs are `dk2nu` and `dkmeta` objects [77] for each neutrino event returning a set of correction values to be used as weights to calculate the right neutrino yield. Internally, PPFX:

- Accounts for the attenuation of all particles passing through the relevant NuMI materials: the target and Budal monitors, the magnetic horn inner conductors, the decay pipe volume and the decay pipe walls.
- Implements energy scaled thin target data for incident particles in the energy

⁸The uncertainties associated with the focusing process have not been implemented yet and are currently handled inside the MINERvA framework.

⁹An extension to other hadronic models is pending but the prescription followed for the current model can be applied directly.

range $[12, 120]$ GeV. All data is carefully interpolated to smooth transition between bins.

- Implements MIPP NuMI target data for pions and high energy kaons.
- When there is no data, **PPFX** attempts first to use theoretically guided extensions and when it is not possible, uses a data driven best guess.
- Handles correlated and uncorrelated uncertainties using the multi-universe technique for uncertainty propagation.

Figure 4.46 shows a schematic view of how the main classes of **PPFX** work. They are described below:




FIG. 4.46: PPFX flowchart.

4.8.1 MakeReweight

MakeReweight is a singleton class that controls the process. It has to be instantiated and initialized by the user selecting options that are entered in a xml file through the class member **SetOptions**. Three options are currently settable:

- The process mode. Two modes are implemented: **MIPPNuMION** and **MIPPNuMIOff**. When **MIPPNuMION** is selected, the code tries to use thick target data as a primary correction and thin target data for any interaction not yet covered. When **MIPPNuMIOff** is selected, the code do not use thick target at all and apply a correction based on thin target data for all interactions (This switching mechanism will be use for Gen2-thin and Gen2-thick flux versions as will be described in section 5.3).
- The number of universes to be used in the process.
- The MIPP NuMI hadron production bin to bin correlation for systematic uncertainties. The default value is +75%.

MakeReweight is also in charge of initializing:

- The **CentralValueAndUncertainties** class that creates tables of central values and deviates per universe (it is called **ParameterTable**). This is done by reading the input data from a xml file (**parameter file**).
- The binning convention of the data input matching the parameter file convention.
- A vector of **ReweighterDriver**'s (where every entry corresponds to one universe and becomes its identification, ID) and a **ReweighterDriver** for the central value (which has an ID equals -1).

Its member **CalculateWeights** constructs an **InteractionChainData** object using the dk2nu and dkmata inputs and passes them to every **ReweighterDriver** to

make the calculations and provides to the user a vector of universe weights by calling the member `GetWeights()` and a central value weight by `GetCVWeight()`.

4.8.2 Data and MC Inputs

The parameter file contains the input data from results of studies to make interpolations and correlations between bins. The implementation of every dataset has been presented and discussed during the MINERvA flux group meetings (which can be tracked looking at MINERvA doc-db ¹⁰). For uncorrelated data, the central value and its uncertainty are provided as single values or a list of values. Correlated data requires a list of central values per bin, a list of statistical errors and a covariance matrix of the systematic errors. The `CentralValueAndUncertainties` class parses the parameter file and its member `calculateParsForUniverse` calculates a parameter table with all random data values per universe, where every universe has a unique and fixed random seed to make the process replicable and stable.

The MC values have been tabulated offline and, except for MIPP NuMI that enters as an xml file ¹¹, they are incorporated as root histograms. Three classes store these values and provide them as request by reweighters: `AttenuationMC`, `ThinTargetMC` and `MIPPNumiMC`.

4.8.3 InteractionChainData

The `InteractionChainData` class parses `dk2nu` and `dkmeta` objects and classifies the information in:

- A vector of `InteractionData` objects that contains information for an interaction (kinematics, identity of the projectile, target and produced particles).

¹⁰Some useful presentations are [78], [79] and [80] as well the implementation parts of sections 4.4.1 to 4.7.

¹¹This is due to an early implementation before deciding to use root files for MC inputs.

- A `TargetData` object with information for the hadron that exits the target: kinematics and identity of the hadron. This also includes its position index in the vector of `InteractionChainData`.
- A vector of `ParticlesThroughVolumesData` objects that stores the momentum and the amount of material crossed by the neutrino parent, grand-parent and when it exists, its great-grand-parent per volume.

4.8.4 ReweightDriver

The `ReweightDriver` class is in charge of initializing the specialized reweighters for specific interactions and using them to calculate hadron production weights. A reweighter makes the task for one universe or for the central value receiving the corresponding `ParameterTable` and the `InteractionChainData` from the `MakeReweight` class. The member `CalculateWeights` selects the appropriate reweighter and book-keeps a history of the interactions already corrected to avoid duplications.

4.8.5 Reweighters

These classes are responsible for the hard work, i.e. calculating the weights. Those are inherited from one of the two following abstract base classes:

- `IInteractionChainReweighting`. The interface for classes that correct a chain of interactions given an `InteractionChainData`. Two members have to be implemented by reweighters inherited by this class: `canReweight` and `calculateWeight`. The former looks at the `InteractionChainData` input and identifies those interactions that can be corrected as part of a chain and returns a vector indicating which elements will be assigned a weight by `calculateWeight`.

- **IInteractionReweighting**. The interface for classes that corrects one interaction given by **InteractionData**. As before, **canReweight** and **calculateWeight** have to be implemented by every reweighter.

The reweighters associated with the attenuation correction are special because even when they are inherited from **IInteractionChainReweighting**, they are not specialized in correcting hadron production interactions and they use the **InteractionChainData** to get the vector of **ParticlesThroughVolumesData**.

The rest of the section is devoted give a brief description of the data used and their implementation in reweighter.

The complete list of reweighters is:

Type	Name	Sections
Thick target	MIPPNumiPionYieldsReweigher	4.4.1
	MIPPNumiKaonYieldsReweigher	4.4.2
Thin target	ThinTargetpCPionReweigher	4.5.4
	ThinTargetpCKaonReweigher	4.5.5
	ThinTargetpCNucleonReweigher	4.5.6
	ThinTargetnCPionReweigher	4.6.1
	ThinTargetnucleonAReweigher	4.6.1
	ThinTargetMesonIncidentReweigher	4.6.2
	OtherReweigher	4.6.2
Attenuation	TargetAttenuationReweigher	4.7.1 - 4.7.3
	AbsorptionICReweigher	
	AbsorptionDVOLReweigher	
	AbsorptionDPIPReweigher	
	NucleonAbsorptionOutOfTargetReweigher	
	OtherAbsorptionOutOfTargetReweigher	

TABLE 4.4: PPFX reweighters.

CHAPTER 5

A priori MINERvA Low Energy Beam results

This chapter is dedicated to presenting the *a priori* LE flux results at MINERvA. As was mentioned in section 3.5, the flux prediction has been permanently updated to add new hadron production data, increase the accuracy of the NuMI simulation's geometry and improve the treatment of the flux systematics.

Historically, we have had two NuMI flux predictions called Generation 0 and Generation 1 that were used in the current published MINERvA papers. Table 5.1 has a list of the MINERvA papers for each flux generation. The most updated version is Generation 2 which includes the hadron production data constraints described throughout sections 4.4 to 4.7.

There are some commonalities between flux generations:

- The use of the multi-universe technique (see section 4.2.5) to propagate the hadron production and focusing uncertainties.
- The use of G4numi as MC flux simulation and FTFP_BERT as the hadronic physics model.

Additionally, it is worthwhile to emphasize that the MINERvA flux results are calculated integrating over the MINERvA fiducial volume defined as the region inside of an hexagon with 86 cm apothem transverse area to the beam axis and within a 126.7 cm region along z axis corresponding to the “downstream” portion of tracker region. Our procedure is the following:

- We select neutrino inclusive charged-current interactions on carbon produced in the fiducial volume. The selection is made using the MC truth information.
- Based on the neutrino ancestry record, we apply hadron production corrections and propagate the uncertainties associated to the hadron production and the focusing system. For Generation 0 and Generation 1, this is made using a code inside the MINERvA offline computational framework. For Generation 2, we use PPFX that is plugged into the MINERvA framework as an external package.
- The event rate per neutrino type is then divided by the corresponding GENIE cross-section and the number of carbon atoms in the fiducial volume.

This chapter is structured in 4 sections. In the first two sections, we will explain the procedure and results for Generation 0 and Generation 1. We present the Generation 2 results in section 5.3 offering also an analysis of the contribution of every PPFX reweighter. In the final section, we discuss the implications of the Generation 2 results.

5.1 MINERvA flux Generation 0

Generation 0 has been calculated with the following components:

- **Hadron production central value.** We only used thin target data to correct interactions in the target. The energy scaling described in section 4.2.2 is also used. We applied:

- NA49 (section 4.5.1) and Barton (section 4.5.2) charged pion production on proton - carbon interactions (figure 4.25).
- NA49 charged kaon production on proton - carbon interactions (figure 4.29).
- NA49 proton production on proton - carbon interactions (figure 4.36).
- Attenuation of the primary protons that interact in the target were applied using NA49 inelastic cross-section (sections 4.2.1 and 4.7.1).

- **Hadron production uncertainties.** The uncertainties applied were:

- The statistical and systematic uncertainties added in quadrature of the NA49 pion production on proton - carbon interactions to the primary proton interactions producing pions in the target (figure 4.27 and table 4.2).
- The model spread between geant4 hadronic models was applied to the rest of the interactions 4.2.4.

- **Focusing:**

- We applied the focusing uncertainties inherited from MINOS [20]: transverse horn offset (section 2.5.1), baffle scraping (section 2.5.2), POT counting (section 2.5.3), horn current magnitude (section 2.5.4), current distribution in the horn inner conductors (explain in section 2.5.4) and the longitudinal target offset (section 2.5.6).
- The flux is simulated using the nominal target positions (table 2.3), i.e., it was before of the longitudinal target position survey described in section 2.5.6.

Figure 5.1 a) shows the Generation 0 ν_μ and $\bar{\nu}_\mu$ fluxes from LE010z185i and LE010z-185i respectively before the HP correction and after applying the correction. The size and energy dependence of the HP correction can be seen in 5.1 b).





FIG. 5.1: ν_μ and $\bar{\nu}_\mu$ Generation 0 flux.

5.2 MINERvA flux Generation 1

Generation 1 has been calculated considering:

- **Hadron production Central value.** In addition to the datasets included in Generation 0, we used:
 - MIPP K/π yield ratios to extend the NA49 data coverage (figure 4.30).
 - We apply the HP correcting to all materials (not only to the target) using the same weights calculated for carbon. This was the first effort to extend the data coverage to other materials but without any scaling studies yet.
 - We correct a bug on our computation of the MC invariant cross-section for π^\pm . The Geant4 hadronic models produce explicitly η , η' , Σ^0 and π^0 when simulating the hadronic interactions. Hadron production experiments do not explicitly measure η and η' and they can decay into charged pions. Then, cases in which there is a charged pion from an $\eta(\eta')$ decay has to be accounted when calculating the invariant cross sections.
- **Hadron production Uncertainties.** There was no change in the hadron production uncertainties.
- **Focusing.** There was no change in the focusing uncertainties.

Figure 5.2a shows the Generation 1 ν_μ and ν_e fluxes from LE010z185i, respectively, and $\bar{\nu}_\mu$ flux from LE010z-185i before the HP correction and after applying the correction. The size and energy dependence of the HP correction can be seen in 5.2b. The differences with Generation 0 around the focusing peak come mainly from the fix of the $\eta(\eta')$ bug.

Figure 5.3 shows the ν_μ and ν_e flux fractional uncertainties ($\bar{\nu}_\mu$ is similar to ν_μ). As was explained before, the uncertainties did not change when we moved





FIG. 5.2: Generation 1 flux.

from Generation 0 to Generation 1. The uncertainty from the model spread (“no data constraint” in the plot) dominates the ν_μ total uncertainty except around the focusing peak where NA49 data is used for most of the interactions that contributes to that neutrino energy region (red line). For ν_e flux, the large contribution of highly energetic kaon neutrino parents makes the model spread (and the total uncertainty) be higher than ν_μ .

Table 5.1 contains a complete list of the MINERvA papers currently published. As was mentioned before, they will need to be updated with Generation 2 flux results (see chapter 6).

Flux	Analysis
Generation 0	ν_μ Charged Current Quasi-elastic Scattering [81]
	$\bar{\nu}_\mu$ Charged Current Quasi-elastic Scattering [82]
Generation 1	Charged Current Scattering Target Ratios [83]
	Coherent π Production [5]
	Muon + Proton Production in ν_μ Scattering [84]
	Charged Current π^0 Production in $\bar{\nu}_\mu$ Scattering [85]
	Charged pion production in ν_μ interactions [86]


TABLE 5.1: Flux used in the current MINERvA publications.

5.3 MINERvA flux Generation 2


In may of 2014, we started working on a new flux version - Generation 2 - motivated by:

- **Hadron Production:**

- Thick target data was published by the MIPP experiment using a LE NuMI target (see section 4.4.1).
- The need to reduce the flux uncertainties and gain a deeper understanding of them by replacing the model spread with direct data or physics motivated data extension corrections (section 4.2.4).



(a) ν_μ .



(b) ν_e .

FIG. 5.3: Generation 0 and Generation 1 flux fractional uncertainties for ν_μ and ν_e from LE010z185i.

- The inclusion of particle absorption in the beamline volumes (section 4.2.1) beyond primary proton in the target.

- **Improvements in the geometry simulation of NuMI that implement:**

- an accurate target positions (section 2.5.6).
- a water layer around the horn inner conductors and other missing materials (section 2.5.7).
- an improved horn geometry model (section 2.5.5).

In this section, we present the Generation 2 Low Energy Beam flux results for ν_μ in minerva13, $\bar{\nu}_\mu$ in minerva5 and ν_e in minerva13 (see section 2.5.6 for the playlist definition).

The implementation of thin and thick target data in PPFX, following the procedure in section 4.2, gives us the opportunity to have two flux versions that are called Generation2 - thin (**Gen2-thin**) and Generation2 - thick (**Gen2-thick**).

Gen2-thin uses primarily thin target data to correct hadronic interactions (see section 4.5). The first attempt is to use data directly and then, when that is not possible, we look to extend the data (i.e. for pA and nA interactions, see section 4.2.4). For the remaining interactions ($\pi A \rightarrow \pi A$, etc.), we make an educated guess (see section 4.2.4). The attenuation of the particle beams is corrected for any hadron passing through the relevant NuMI volumes: TGT, IC, DVOL and DPIP (see section 4.7).

Figures 5.4 a) and b) show the average number of hadronic interactions per ν_μ and ν_e in MINERvA that are handled by each reweighter in **Gen2-thin** (the $\bar{\nu}_\mu$ plots are not shown here since are similar to the ν_μ ones). The black lines represent the total average. Except for low neutrino energies (< 1 GeV), these values are around 1.4, meaning that a large fraction of these interactions happen inside of the

target as we can see when we compare this with the interactions covered by hadrons that exit the target (figures 4.12 and 4.13).

Every line represents the coverage of one PPFX reweighter (see table 4.3):

- MIPP NuMI π and K production are shown by solid and dashed red lines respectively .
- Pion, kaon and nucleon production in proton - carbon interactions using thin target data are shown by solid red, dashed red and dashed orange lines respectively.
- Pion production in neutron carbon interactions constraint with data using isoscalar principles are shown by solid orange lines.
- Meson incident interactions are shown by solid brown lines.
- Nucleon incident interactions not covered directly by any data (it is called **nucleon-A** in the plots) are shown by dashed brown lines.
- **Others** includes the remaining interactions not covered yet ($pC \rightarrow \pi^+ X$ at $x_F > 0.5$ and $p_T > 0.5$ that are not covered by neither NA49 nor Barton).

The dominant corrections handled by PPFX for ν_μ (ν_e) are for pion production from proton carbon interactions up to 20 GeV (8 GeV) after which the kaon production correction becomes relevant at high energies in accordance with their ancestry (see figure 2.16a). The coverage around the focusing peak is almost complete (compare with the figure 2.17¹).

The other significant component is nucleon-A, especially around the transition between pion and kaon production. This is because only a small region with $x_F > 0.5$ is covered by Barton. The nucleon interactions, nucleon production and meson

¹Figure 2.17 shows the average number of interactions integrated over the neutrino energy in 0-20 GeV but it is a good approximation for the focusing peak.

incident corrections have higher values at low energies because they are composed by secondary and tertiary hadron production.

Gen2-thick uses thick target data primarily, and then, for the interactions still not covered, uses thin target data and an educated guess. The beam particle attenuation is corrected for any hadron passing through the IC, DVOL and DPIP. For TGT, only the primary proton beam attenuation (in combination with thick target data) is corrected (see section 4.2.2).

Figures 5.5 a) and b) show the effect of thick target data on the average number of interactions covered by every reweighter in **Gen2-thick**. The lines for MIPP NuMI pion and kaon cover many of these interactions especially around the focusing peak. Thin target pion production becomes relevant only for low energies due to the MIPP acceptance (see 4.15 for an example). The contribution from nucleon-A also decreases, particularly due to the coverage of secondary protons reinteracting in the target to make pions. The small bump for thin target kaon production around 8-19 GeV comes from kaons below 20 GeV covered by NA49 ($x_F < 0.2$ GeV).

I split nucleon-A into more categories in order to understand which corrections are being handled for this reweighter: figure 5.6 a) for **Gen2-thin** and 5.6 b) for **Gen2-thick**. Some of the characteristics are:

- Pion and kaon production from nucleon interactions (solid blue line) using the material scaling procedure described in section 4.2.3. This value is small in the focusing peak because more of the neutrino parents are produced in the target. For higher energies it is almost flat (~ 0.11) due to primary and high x_F secondary protons interacting outside the target. This line remains unchanged between **Gen2-thin** and **Gen2-thick** as expected.
- The quasi-elastic proton production outside of the target (dashed red line) remain unchanged as expected and it is a very small component of nucleon-A.





FIG. 5.4: **Gen2-thin** average number of hadronic interactions per neutrino minerva13





FIG. 5.5: **Gen2-thick** average number of hadronic interactions per neutrino minerva13

- The incident nucleon with momentum less than 12 GeV (dashed blue line) has a small contribution to nucleon-A and remain almost unchanged since or it is not covered by thin target data or it produces low energy pions in the not MIPP acceptance region.
- Some of the protons created in quasi-elastic interactions in the target are also going to interact in the target and produce pions or kaons covered by MIPP. That is why this component (solid red line) reduces its value by more than half in **Gen2-thick** respect to **Gen2-thin**.
- The orange line represents the remaining nucleon production after applying the previous classification.
- The category **others** represents any nucleon-A interaction not classified yet. As was said, it has a significant value at the region between the edge of the pion production and the rise of kaon production PPFX coverage (neutrino energies in 20-30 GeV). They are basically pions or kaons not covered by data (directly or by using the material scaling) and produced in any material (for **Gen2-thin**) or just outside of the target (for **Gen2-thick**).

Figure 5.7 a) shows the average material traversed by any hadron in the relevant NuMI volumes for the beam attenuation effect: TGT, IC, DVOL and DPIP per ν_μ with energies between 0-40 GeV that passes through MINERvA detector. **Gen2-thin** corrects the beam particle attenuation for the total average material traversed (black line) shown in the plot while **Gen2-thick** uses the primary proton passes through carbon (solid green line) and all lines in materials different than carbon. The same information but just for pion neutrino parents is in figure 5.7 restricted to the neutrino energy 0-20 GeV. Some characteristics to highlight are:

- For almost the whole neutrino energy range, the primary proton beam traversed





FIG. 5.6: Nucleon-A interaction per neutrino covered in Generation 2.

about 40 cm ($\sim 6 \text{ mol/cm}^2$) in the target as average, except for neutrino energies around the focusing peak where the pions produced in the primary proton interactions are focused optimally: the average distance traveled by the primary protons decreases from $\sim 27 \text{ cm}$ to $\sim 17 \text{ cm}$ while for pions increases from $\sim 7 \text{ cm}$ to $\sim 17 \text{ cm}$ when the neutrino energy goes from 2 GeV to 5 GeV.

- The pions traverse about 10 cm ($\sim 1 \text{ mol/cm}^2$) of Al on average. This is because the magnetic horns are long and thin and the pion angles are small and tend to travel more distance inside the horn inner conductors.
- Pion neutrino parents with energies greater than $\sim 4.5 \text{ GeV}$ (using the equation 2.1) tend to reach the decay pipe and travel $\sim 200 \text{ m}$ before decaying.
- As the neutrino energy increases, the material traversed by kaon becomes relevant.

5.3.1 Results

This subsection shows the flux results for:

- **Gen2-thin**, ν_μ , figure 5.8,
- **Gen2-thin**, $\bar{\nu}_\mu$, figure 5.10,
- **Gen2-thin**, ν_e , figure 5.12,
- **Gen2-thick**, ν_μ , figure 5.9,
- **Gen2-thick**, $\bar{\nu}_\mu$, figure 5.11,
- **Gen2-thick**, ν_e , figure 5.13.





FIG. 5.7: NuMI material traversed.

The energy range is restricted to 0-20 GeV. The error bands combine hadron production and focusing uncertainties. The upper plots show the flux spectrum and the lower plot the ratio to the g4numi output in order to indicate the final correction we applied after using the tools developed in **PPFX**. A complete table, bin by bin, can be found in appendix B.

For ν_μ , the MC tends to underestimate the number of neutrino below 8 GeV for **Gen2-thin** and below 11 GeV for **Gen2-thick** (except at very low energies) and overestimate everywhere else. A discussion of the agreement between the two predictions can be found in section 5.4, but it is clear that **Gen2-thick** predicts a big "dip" around 4-10 GeV of more than 20% while this effect in **Gen2-thin** is not that high.

The hadron production uncertainties for ν_μ and ν_e in **Gen2-thin** are shown in figure 5.14 ($\bar{\nu}_\mu$ plots are not shown since they are similar to ν_μ) for an energy range of 0-20 GeV.

For **Gen2-thin**, the total uncertainties around the focusing peak are approximately 6-7% for both ν_μ and ν_e due to the NA49 pion production data coverage. The uncertainties for ν_μ increase to $\sim 9\%$ for energies beyond 6 GeV due the influence of the nucleon-A correction, however, ν_e uncertainties increases in the same region because the kaon production becomes relevant. In general the uncertainties are below 4% for the interactions where we applied data directly. The attenuation in TGT and other materials (it is called "absorption" in plots) has an uncertainty of $\sim 3\%$.

For **Gen2-thick** (see figure 5.15), the total uncertainties are even smaller with an almost flat value about $\sim 5\%$. This is due to the MIPP NuMI data coverage. All the rest of the components reduce their value, except "absorption", since that is applied to volumes outside of the target.




FIG. 5.8: Gen2-thin ν_μ flux.




FIG. 5.9: Gen2-thick ν_μ flux.




FIG. 5.10: Gen2-thin $\bar{\nu}_\mu$ flux.




FIG. 5.11: Gen2-thick $\bar{\nu}_\mu$ flux.




FIG. 5.12: Gen2-thin ν_e flux.




FIG. 5.13: **Gen2-thick** ν_e flux.





FIG. 5.14: Fractional uncertainties for Gen2-thin.







FIG. 5.15: Fractional uncertainties for Gen2-thick.

The bin to bin energy correlation between ν_μ , $\bar{\nu}_\mu$ and ν_e is shown in figure 5.16 for **Gen2-thin** and **Gen2-thick**. Some observations can be made from these plots:

- bins with high correlation within each neutrino flavor reflect the highly correlated systematic uncertainties assumption we have made for thin and thick target data and the contribution of different data bins to every neutrino energy bin.
- the correlation in the transition from lower to higher energies around 5-6 GeV is due to the focusing uncertainties.
- The correlation between ν_μ and $\bar{\nu}_\mu$ comes from neutrinos with more than one hadronic interaction in their cascade history that share the same interaction for one or more ancestries. Comparing with Gen2-thick, when the ν_μ and $\bar{\nu}_\mu$ correlation is very weak, we can conclude that these common interactions mainly happen inside of the target.



(a) Gen2 - thin



(b) Gen2 - thick

FIG. 5.16: Energy bin to bin flux correlation.

5.4 Discussion

The flux integrated values over 0-20 GeV are shown in table 5.2 for ν_μ , $\bar{\nu}_\mu$ and ν_e . **Gen2-thin** and **Gen2-thick** agree at the 1 sigma level. The uncertainties in the integrated flux have similar values of the flux spectrum around the focusing peak as we expect. The correlations between ν_μ and $\bar{\nu}_\mu$ are shown in figures 5.17 and 5.18.

	integral ($\nu/m^2/10^6\text{POT}$)	Uncertainty (%)
Gen2-thin ν_μ	287.0	7.78
Gen2-thick ν_μ	280.8	5.37
Gen2-thin $\bar{\nu}_\mu$	233.5	7.46
Gen2-thick $\bar{\nu}_\mu$	238.6	5.51
Gen2-thin ν_e	4.11	7.06
Gen2-thick ν_e	4.07	4.93

TABLE 5.2: Integrated flux for ν_μ and $\bar{\nu}_\mu$ in **Gen2-thin** and **Gen2-thick** for neutrino energy in 0-20 GeV.




FIG. 5.17: **Gen2-thin** flux integral correlation between ν_μ and $\bar{\nu}_\mu$. Each entry in this histogram is one universe.

The ratio of **Gen2-thick** over **Gen2-thin** flux spectra in figures 5.19 and 5.20 show the level of consistency between both flux predictions. The ratios have




FIG. 5.18: **Gen2-thick** flux integral correlation between ν_μ and $\bar{\nu}_\mu$. Each entry in this histogram is one universe.

been made canceling all possible correlations to be sensitive to the real size of any disagreement between both versions. We expect that if they agree perfectly, the ratios should be compatible with 1, within errors.

For $\bar{\nu}_\mu$, both predictions agrees within errors. However, for ν_μ , there is a energy dependent disagreement that is greater than 1 sigma in two energy regions: 5-13 GeV and 20-35 GeV. The former region is crucial to understand since it covers the falling edge of the focusing peak and most of the MINERvA analysis are restricted to energies below 20 GeV. Table 5.20 shows the bins when the disagreement is more than 1σ for ν_μ below 30 GeV. This conflict will be explored and resolved in the next chapter.

$\Delta E(GeV)$	5-6	6-7	7-8	8-9	9-10	10-11	11-12	12-13
σ	1.52	1.97	2.16	2.04	1.96	1.90	1.86	1.44

TABLE 5.3: Disagreement between **Gen2-thin** and **Gen2-thick** for ν_μ .




FIG. 5.19: ν_μ Gen2-thin and Gen2-thick comparison.




FIG. 5.20: $\bar{\nu}_\mu$ Gen2-thin and Gen2-thick comparison.

CHAPTER 6

Additional flux constraining using MINERvA in-situ measurements

At the end of the last chapter, we showed a potential disagreement between the **Gen2-thin** and **Gen2-thick** the ν_μ flux predictions . Is it possible to reconcile these two predictions, or favor one of them in a physics motivated way? The first section of this chapter addresses this question.

The strongly energy dependent ratio between **Gen2-thick** and **Gen2-thin** ν_μ fluxes (figure 5.19) suggests that the best approach is to use the flux predicted from low-nu since it is essentially a flux shape measurement. Furthermore, as we saw in table 5.2 shows, the flux integrated in all neutrino types agree each with other to within their uncertainties meaning the neutrino - electron scattering constraint does not suitably solve this potential disagreement. However, the second section of this chapter is dedicated to exploring the application of the MINERvA results of this channel to reduce the flux uncertainties.

6.1 Low-nu flux at MINERvA

The low-nu procedure was applied in MINERvA to calculate the ν_μ and $\bar{\nu}_\mu$ fluxes and these fluxes in turn were used to extract inclusive scattering cross-sections [43]. The general principle of the method was explained in section 3.3 and this section presents only the results and a brief summary of the procedure in order to make the comparisons with Generation 2 *a priori* results introduced in the previous chapter. More details can be found in [43].

The low-nu flux analysis was done using the Generation 1 flux prediction, but it is insensitive to the flux model, as one would hope. It uses three ν cuts that were applied to the inclusive scattering events in overlapping neutrino energies: $\nu < 300 \text{ MeV}$ ($E > 2 \text{ GeV}$), $\nu < 800 \text{ MeV}$ ($E > 5 \text{ GeV}$) and $\nu < 2 \text{ GeV}$ ($E > 9 \text{ GeV}$). Figure 6.1 a) shows the fraction of the inclusive data sample (overlap) with less ν less than the cut for ν_μ in LE010z185i¹.

After applying a Bayesian unfolding to account for the detector resolution and after applying efficiency corrections, the three event rate samples under the ν cuts have, basically, the shape of the flux. The next steps followed were:

1. **ν cross-section (σ_ν).** The neutrino nucleon charged current cross-section σ_ν is calculated independently in each three ν samples using equation 1.3 and the *a priori* Generation 1 flux. Figure 6.1 b) shows these cross-sections for ν_μ .
2. **Finite ν .** A correction for finite ν is made using GENIE by calculating:

$$S(\nu_0, E) = \frac{\sigma(\nu < \nu_0, E)}{\sigma(\nu < \nu_0, E \rightarrow \infty)} \quad (6.1)$$

where ν_0 is a particular ν cut and E is the neutrino energy (see equation 3.1).

¹The plots corresponding to figures 6.1 and 6.2 for $\bar{\nu}_\mu$ can be found in [43].





FIG. 6.1: Overlap and extracted ν cross-sections for ν_μ in LE010z185i (taken from [43]).

3. **Normalization.** The extracted charged current cross-section is normalized to match the GENIE cross-section value in the 9 – 12 GeV energy region ² and other normalization factors for all three ν samples are also calculated to agree with this match by following a fitting procedure between ν samples (fit errors are included as uncertainties).
4. **Flux.** The ν_μ and $\bar{\nu}_\mu$ fluxes are extracted by applying the factors founded in step 3 per ν sample and they are merged to get a flux determination in the 2-50 neutrino energy region in 15 bins. Table 6.1 contains a list of the bin convention used in this analysis.

Bin	range (GeV)	Bin	range (GeV)	Bin	range (GeV)
1	2-3	6	7-9	11	22-26
2	3-4	7	9-12	12	26-30
3	4-5	8	12-15	13	30-36
4	5-6	9	15-18	14	36-42
5	6-7	10	18-22	15	42-50

TABLE 6.1: Low-nu binning convention.

The flux results following this procedure (for brevity we called it “low-nu flux”) are shown in figure 6.2 a) for ν_μ in the LE010z185i configuration ($\bar{\nu}_\mu$ flux can be found in [43]). The spectrum is shown at the right and the data/MC ratio on the left. The black markers in the plots correspond to the extracted low-nu flux and the red line is the Generation 1 flux prediction. The total fractional uncertainty, shown in figure 6.2b, is between 8-11%. The contribution from the *a priori* flux to the uncertainty in this calculation is small.

²This step requires to use precised external measurements. This analysis used GENIE since this is itself based on data.





FIG. 6.2: ν_μ flux and its fractional uncertainties in LE010z185i calculated using low-nu technique in MINERvA (taken from [43]).

6.1.1 Generation 2 vs low-nu flux comparison

Figures 6.3 and 6.4 show a comparison between the Generation 2 fluxes (thick and thin) to the low-nu flux for ν_μ . In the ratio plots (lower side), the uncertainties have been computed by adding in quadrature the Generation 2 and low-nu uncertainties³. In principle, adding in quadrature is wrong as it does not account for correlations due to the flux model. However, the flux uncertainty component of the low-nu results is negligible, so the overestimation of these ratio uncertainties is small.

A first “by eye” conclusion is that both, **Gen2-thick** and **Gen2-thin**, agree with the low-nu around the focusing peak and for high energies but **Gen2-thick** disagrees with low-nu in the 5-15 GeV neutrino energy while **Gen2-thin** does not. Since the neutrino energy bins are highly correlated (see figure 5.16), a more careful approach is needed. MINERvA conducted a quantitative study using a χ^2 test [87] fully including the correlations between both fluxes.


The idea is to test the hypothesis (H_0) that $L - G = 0$, where G and L are the Generation 2 (thin or thick) and the low-nu fluxes. A χ^2 is constructed by:

$$\chi^2 = \sum_{i,j;i \leq j}^N (G_i - L_i)(G_j - L_j)V_{ij}^{-1} \quad (6.2)$$


Where the sum is made bin by bin (i,j) for $G_{i,j} - L_{i,j}$ and the matrix $V = V_G + V_L$ is the sum of the Generation 2 and low-nu covariance matrices. The results for two neutrino energy ranges, 2-50 GeV and 2-22 GeV, are summarized in table 6.2. Looking at both ranges, we conclude that **Gen2-thin** is in a good agreement with low-nu while **Gen2-thick** has poor agreement (especially when we consider the whole energy region).

Similar work has been conducted in [87] to test the agreement between **Gen2-**

³Currently, we do not have the technology to cancel common correlations.




(a) Flux spectra




(b) Ratios

FIG. 6.3: Gen2-thin ν_μ flux vs low-nu



(a) Flux spectra



(b) Ratios

FIG. 6.4: Gen2-thick ν_μ flux vs low-nu

Comparison	2-50 GeV	2-22 GeV
	χ^2/NDF	χ^2/NDF
using full covariance matrix		
$G^{thin} - L$	7.3/15	4.8/10
$G^{thick} - L$	61.3/15	18.6/10
using diagonal elements only		
$G^{thin} - L$	8.2/15	5.3/10
$G^{thick} - L$	68.5/15	38.3/10

TABLE 6.2: χ^2/NDF for Generation 2 and low-nu difference. Values taken from [87].

thick and **Gen2-thin** adopting the low-nu binning described in table 6.1. The χ^2 test of the hypothesis $G^{thick} - G^{thin} = 0$ has been calculated. Four 0.5 GeV size bins have been added to cover energies less than 2 GeV. The results are shown in table 6.3. The agreement between both fluxes is poor in all cases. However, this results strongly depends of the correlation assumptions we have made when incorporating the data. One possible solution would be to study the effect of different correlation assumptions in order to find a way to “reconciliate” both flux predictions. This work can be done especially in **Gen2-thin**, where we assume 100% bin-to-bin correlation for NA49 data. Given that some of the data uncertainties, like the normalization, have definitive positive correlation and the big discrepancy in the χ^2 study, reconciling both predictions, **Gen2-thick** and **Gen2-thin**, would have to take not well-founded assumptions, like strong negative bin-to-bin correlations for some systematics. For these reasons consider **Gen2-thin** and **Gen2-thick** to be in disagreement.

	0-50 GeV	2-50 GeV	2-22 GeV
	χ^2/NDF	χ^2/NDF	χ^2/NDF
full covariance matrix			
	144.7/19	91.0/15	76.1/10
diagonal elements only			
	62.1/19	25.4/15	22.0/10

TABLE 6.3: χ^2/NDF for **Gen2-thick** and **Gen2-thin** difference. Values taken from [87].

Summing up this section: Generation 2 gives us the chance to have two flux predictions. Nevertheless, the studies presented suggest that both predictions are in disagreement each other and only **Gen2-thin** is in agreement with the low-nu flux results while **Gen2-thick** does not. Then, **Gen2-thin** will be used as the flux determined by this thesis.

6.2 Additional constraint using neutrino electron scattering events

An additional constraint based on the neutrino electron scattering events can be applied to the flux, on top of the hadron production corrections. As was mentioned in section 3.4 the cross-section for this reaction is well determined by the Standard Model since all participants are leptons.

Figure 6.5 shows the energy distribution of the neutrino-electron elastic scattering candidates interacting in the MINERvA tracker. In total, MINERvA isolated 135 ± 17 ($\sim 12.6\%$) after the background subtraction and efficiency correction. These events were collected using 3.43×10^{20} POT in LE010z185i.

We implement the $\nu e \rightarrow \nu e$ constraint by comparing the predicted number of these events from the *a priori* flux with the measured number. Then, we use this comparison to weight up or down the more likely or unlikely universes by computing the likelihood \mathcal{L}_i for each universe i:

$$\mathcal{L}_i(\vec{N}_{\nu e \rightarrow \nu e} | \vec{Q}) = \frac{1}{(2\pi)^{M/2} \det(V)^{1/2}} \exp\left(-\frac{1}{2}(\vec{N} - \vec{Q}) \cdot \mathbf{V}_N^{-1} \cdot (\vec{N} - \vec{Q})\right) \quad (6.3)$$

where M is the number of data bins. \vec{N} and \vec{Q} are vectors of the measured number and the predicted number, respectively. Every entry in those vectors is the




FIG. 6.5: Electron energy distribution from ν -e scattering measured by MINERvA after the background subtraction and efficiency correction (taken from [88]).

corresponding bin content. \mathbf{V} is the data covariance matrix. These likelihoods are stored and can be used as weights that multiplies any quantity that varies with the flux, particularly, the flux itself.

Figure 6.6a shows how the predicted distribution of $\nu e \rightarrow \nu e$ from the *a priori* Generation 1 flux changes after applying this constraint. Every entry in the *a priori* plot (black line) is the corresponding value in a universe i and, it is multiplied by \mathcal{L}_i and renormalized to preserve the original number of universes (red line). The mean shift and the uncertainty is reduced significantly for more than 30%.




FIG. 6.6: Predicted $\nu e \rightarrow \nu e$ events from the *a priori* flux before and after the constraint (taken from [88]).

The weights from this constraint have been calculated and apply to Generation 2 in MINERvA [89]. Figure 6.7a shows the effect on the ν_μ Gen2-thin flux. The black lines represent the *a priori* Gen2-thin flux and the red line the effect of applying the $\nu - e$ constraint. This produces a reduction of the flux peak by approximately 2%

and a dip of 4% reduction in the falling edge of the flux. The fractional uncertainty also reduces as can be seen in figure 6.7b.


A measurement of ν_e quasi-elastic and quasi-elastic-like scattering on hydrocarbon was recently published by MINERvA using this procedure on the Generation 1 flux [90]. All future MINERvA publications will likely include this additional constraint.

6.3 Comparison of Gen2-thick with Generation 0 and Generation 1


All MINERvA analyses and results already published by MINERvA need to be updated to the standard flux version, **Gen2-thin** (see a table of the current MINERvA papers: 5.1). This section presents comparisons between earlier versions of the flux, Generation 0 and Generation 1. This provides an idea on how the MINERvA cross-section results would change by **Gen2-thin**. However, other studies needs to be considered in the updating process like the effect of the background when switching flux versions.

This is the list of flux comparisons presented in this sections:

- ν_μ flux **Gen2-thin vs Generation 0** (figure 6.8).
- $\bar{\nu}_\mu$ flux **Gen2-thin vs Generation 0** (figure 6.9).
- ν_μ flux **Gen2-thin vs Generation 1** (figure 6.10).
- $\bar{\nu}_\mu$ flux **Gen2-thin vs Generation 1** (figure 6.11).
- ν_e flux **Gen2-thin vs Generation 1** (figure 6.12).



(a) Flux change



(b) Fractional uncertainty change

FIG. 6.7: Effect of the νe constraint on the Gen2-thin flux. Top: the flux spectrum. Bottom: the fractional uncertainties (taken from [89]). The black line comes from the *a priori* flux and the red line is after applying the $\nu - e$ constraint.

- ν_μ flux **Gen2-thin vs Generation 1 + neutrino electron scattering constraint** (figure 6.13).
- ν_e flux **Gen2-thin vs Generation 1 + neutrino electron scattering constraint** (figure 6.14).
- The flux integrated ratios for ν_μ , $\bar{\nu}_\mu$ and ν_e in the 0-20 energy range (table 6.4).

Neutrino	<u>Gen2-thin</u> <i>Gen0</i>	<u>Gen2-thin</u> <i>Gen1</i>	<u>Gen2-thin</u> <i>Gen1+nu-e</i>
ν_μ	0.846	0.890	0.959
$\bar{\nu}_\mu$	0.838	0.892	-
ν_e	-	0.900	9.970

TABLE 6.4: Integrated flux ratios between generations in the 0-20 energy range. We abbreviate Generation 0 by Gen0, Generation 1 by Gen1 and neutrino electron constraint by nu-e.




FIG. 6.8: ν_μ flux **Gen2-thin** vs Generation 0




FIG. 6.9: $\bar{\nu}_\mu$ flux **Gen2-thin** vs Generation 0




FIG. 6.10: ν_μ flux **Gen2-thin** vs Generation 1




FIG. 6.11: $\bar{\nu}_\mu$ flux **Gen2-thin** vs Generation 1




FIG. 6.12: ν_e flux **Gen2-thin** vs Generation 1




FIG. 6.13: ν_μ flux **Gen2-thin** vs Generation 1 + neutrino electron scattering constraint




FIG. 6.14: ν_e flux **Gen2-thin** vs Generation 1 + neutrino electron scattering constraint

CHAPTER 7

Medium Energy Beam results

The focus of this thesis has been to determine the a priori LE flux for MINERvA experiment. However, the procedure we followed as well as the tools we developed are intended to be directly applicable to calculate the Medium Energy flux for any point in the NuMI beamline. As we mentioned in the introductory chapter (section 2), experiments like MINERvA, MINOS and NOvA are currently taking data for the neutrino cross-section and oscillation physics programs at Fermilab and they can benefit from the results of this thesis.

The objective of this chapter is to show our results for the ME flux at MINERvA (first section) and the capability of our techniques to extend to any detector that sees ν_μ neutrinos (in the second section, we present our flux prediction for NOvA Near Detector as an example). For the results shown in this chapter, we used **Gen2-thin** flux version as it was determined that has better physics consistency than **Gen2-thick** (see section 5.4 and 6.1.1).

7.1 Gen2-thin flux for ME MINERvA

Most of the discussion of the LE flux at MINERvA can be applied to understand the results when we switch to the ME beam mode. In this section, we will emphasize only those that are particular to the ME mode. The simulated ME flux spectrum at MINERvA (before any HP constraint) was shown in figure 2.14 with its characteristic peak at 6 GeV. The key differences in ME with respect to the LE mode comes from the changes in the relative positions between the target and horn (it was lengthened), between horns (it was enlarged) and the length of the target ($\sim 20\%$ longer) as was described in section 2.3. These changes cause the focusing system to enhance higher energy mesons that then decay to give higher energy neutrinos (for instance, π^+ yields shown in the lower right side of figure 2.13). See section A.1 for details of the focusing components.

Figure 7.1a shows the average number of hadronic interactions per ν_μ that passes through the center of the MINERvA tracker, the total number as a black line and those that are handled by every PPFX reweighter listed in table 4.4. The shapes of every line are similar to the LE configuration (figure 5.4), however, there are more interactions in total while every reweighter has more interactions to handle. This is because to that the ME target is longer than the LE one so more interactions happen inside the target. Pions and kaons produced in proton - carbon interactions increase, but also incident nucleons not handled directly by data decrease since fewer protons leave the target.

The average material traversed by particle beams in the relevant NuMI volumes per ν_μ is shown in 7.1 a). It also has the same trend as LE (figure 5.7 a)) but as we expect, we have more contributions from the primary proton beam passing through the target. Pion attenuation in the target becomes dominant in the falling edge of the focusing peak and in general, primary protons, pions, and kaons traverse more

material than in LE.

Figure 7.2 shows the flux spectrum for ν_μ in Medium Energy integrated over the MINERvA fiducial volume (see the introduction to chapter 5) in the energy region 0-20 GeV with the error bars corresponding to the HP uncertainties. At the moment this thesis is being written, the ME focusing uncertainties are being revisited. For this reason, we use the focusing uncertainties corresponding to the LE100z200i plus the horn inner conductor uncertainty (section 2.5.5) calculated for ν_μ . Nevertheless, we expect that this is a good approximation of the final ME focusing uncertainties. A table of the ME flux is shown in table B.3 for all neutrino types.


The ν_μ hadron production fractional uncertainty is shown in figure 7.3. It has a similar shape but slightly larger than LE as we can see when we compare with figure 5.14a. However, as we expect, the absorption and incident meson uncertainties are larger.

The correlation between energy bins is shown in figure 7.4 for ν_μ and ν_e . Only the hadron production uncertainties are considered. The bins are highly correlated due to our assumption about the bin-to-bin correlation of the thin target data systematic uncertainties.


Table 7.1 contains the ν_μ , $\bar{\nu}_\mu$ and ν_e integrated over the 0-20 GeV region with the hadron production uncertainties.

	integral ($\nu/m^2/10^6\text{POT}$)	Uncertainty (%)
Gen2-thin ν_μ	704.6	7.26
Gen2-thin $\bar{\nu}_\mu$	0.39	10.98
Gen2-thin ν_e	6.90	6.76

TABLE 7.1: ν_μ , $\bar{\nu}_\mu$ and ν_e **Gen2-thin** integrated fluxes in 0-20 GeV.



(a) Average number of interactions



(b) Material traversed

FIG. 7.1: ME PPFx coverage for ν_μ in MINERvA.




FIG. 7.2: Medium Energy Gen2 - thin ν_μ flux at MINERvA




FIG. 7.3: Fractional Uncertainties for ME **Gen2-thin** ν_μ flux in MINERvA.




FIG. 7.4: Bin to bin correlation of the ME **Gen2-thin** ν_μ and ν_e fluxes at MINERvA.

7.2 Gen2-thin for Nova ND


As an example of the extension of our procedure to other NuMI detectors, we show the ν_μ **Gen2-thin** flux calculated at the center of the NOvA Near Detector front face. Since the flux spectrum peaks at 2 GeV, we restricted the plots shown in this section to the 0-6 GeV energy region.

Figure 7.5a shows that the average number of hadronic interactions per neutrino is higher than the ME beam at MINERvA due to the off-axis position of the NOvA Near Detector. Most of the interactions are handled by the thin target data, particularly pion production from proton - carbon interactions. However, other interactions become relevant at low energies (< 0.8 GeV), particularly nucleons that interact outside of the target that are not covered by data. Figure 7.5b shows the material traversed by the particles in the beam. The contribution of mesons is almost constant with energy since the effect of the focusing is not as relevant in an off-axis beam with respect to the on-axis beam. The category “others-C” in the plots is composed mainly by secondary protons.


Figure 7.6 shows the ν_μ flux spectrum and the error band from the hadron production uncertainties in 0-6 GeV energy region (table B.4 contains these values in 0.1 GeV bin size). The ratio plot indicates that our NuMI flux simulation (g4numi) overestimates the neutrino flux at NOvA by $\sim 10\%$.

As can be seen in figure 7.7, the hadron production fractional uncertainties are $\sim 8\%$ around the focusing peak. The biggest uncertainties come from incident nucleons interacting outside of the target and incident mesons where we are not applying any data constraint.

An interesting comparison can be made between our flux results and the predicted spectrum from the NOvA flux simulation. NOvA uses FLUGG2008 [91] that uses FLUKA for hadronic interactions and GEANT4 for the detector geometry



(a) Average number of interactions



(b) Material traversed

FIG. 7.5: Average number of interactions and material traverse per ν_μ that pass through NOvA in the ME **Gen2-thin** flux.




FIG. 7.6: Gen2-thin ME ν_μ flux.




FIG. 7.7: Fractional Uncertainties for ME Gen2 - thin ν_μ flux in NOvA.

and particle tracking. Figure 7.8 shows the ratio between **Gen2-thin** (called just “ppfx” in the plot) and the FLUKA2008 flux spectra at NOvA ’s Near Detector (called “flugg” in the plot). The uncertainty on the ratio comes from the uncertainty on the **Gen2-thin** prediction. Around the focusing peak, we predict 5-6% less neutrinos and a 15% dip around 2.5 to 4 GeV. There are two possible sources that could explain these discrepancies. First, a mismodeling in the FLUKA hadronic interactions since NOvA does not do a hadron production correction. Second, differences in the NuMI geometry incorporated in g4numi with respect to FLUGG2008 could cause the peak to shift, and added more material like the water layer around the Horn 1 inner conductor (section 2.5.7) could be responsible for attenuating the flux. However, further investigations are needed to give a conclusive explanation.




FIG. 7.8: Comparison between **Gen2-thin** and FLUGG ν_μ flux NOvA ND. The error bar comes from the **Gen2-thin** hadron production uncertainties.

CHAPTER 8

Conclusions

Determining the neutrino flux with small and well-understood uncertainties is crucial for neutrino - nucleus cross-section experiments such as MINERvA and neutrino oscillation experiments such as MINOS and NOvA (chapter 1). We have shown a new computation of the *a priori* NuMI flux constrained by all relevant hadron production data (chapter 4) and using the current knowledge of the NuMI beamline, that reduces the uncertainties and improves error budget accounting. The MINERvA strategy (chapter 3) also includes the use of *in situ* measurements that provide additional information about the flux and can be applied a top of the hadron production constraint. This *in situ* measurements help us check the consistency of the *a priori* flux (the low-nu method) and to serve as an additional constraint (neutrino - electron scattering) (chapter 6).

We calculated two *a priori* flux versions for the LE beam, **Gen2-thin** and **Gen2-thick**. The comparison of these two versions shows an energy dependence disagreement (chapter 5) for the ν_μ flux in the LE010z185i configuration. A quantitative study found that the **Gen2-thin** and low-nu fluxes are in good agreement, whereas the **Gen2-thick** flux agrees poorly with low-nu (chapter 6). Based on this

fact, we presented **Gen2-thin** as the main flux result of this thesis and recommended use it as standard flux for the current and future MINERvA analyses. We also presented a comparison of the **Gen2-thin** LE flux to previous flux versions, Generation 0 and Generation 1. We also presented the **Gen2-thin** ME flux for MINERvA with 7-8% hadron production uncertainties around 6 GeV (chapter 7).

All techniques and tools developed in this thesis (chapter 4) are intended to be available for any other experiment that sees neutrino from the NuMI beamline. Particularly we introduced a computational tool called **PPFX** that is open and public. As an example of extending our procedure to other detectors, we presented the flux calculated in the center of the front face of the NOvA Near Detector with 7-8% hadron production uncertainty around 2 GeV. The extension to other beamlines is straightforward.

Our work indicates where additional hadron production data is needed in order to further reduce uncertainties for NuMI and LBNF. Particularly, it would be very valuable to measure the invariant differential cross-sections of:

- Proton interactions on a thin target with materials different than carbon.
- Quasi-elastic proton production.
- Incident meson interactions producing pions.

Our calculation of the **Gen2-thick** LE flux also shows that the uncertainties can be reduced significantly by using thick target data, like measurements of particle production on a replica of spare target. In this context, the program of measurements proposed by US-NA61 on thin and thick targets is extremely important.

APPENDIX A

Supplementary material for chapter 2

A.1 Focusing components

This section contains:

- NuMI flux focusing components (figure A.1 in LE010185i at MINERvA.
- Longitudinal vs. transverse momentum yield per $\pi^+ \nu_\mu$ parent:
 - Unfocused (figure A.2).
 - Horn2 only (figure A.3).
 - Horn1 only (figure A.3).
 - Underfocused (figure A.5).
 - Overfocused (figure A.6).
 - Others (figure A.7).




FIG. A.1: Longitudinal vs. transverse momentum yield per $\pi^+ \nu_\mu$ parent

Unfocused




FIG. A.2: Unfocused components of $\pi^+ \nu_\mu$ parent

Horn2 only




FIG. A.3: H2 only components of $\pi^+ \nu_\mu$ parent

Horn1 only




FIG. A.4: Horn1 only components of $\pi^+ \nu_\mu$ parent

Underfocused




FIG. A.5: Underfocused components of $\pi^+ \nu_\mu$ parent

Overfocused




FIG. A.6: Overfocused components of $\pi^+ \nu_\mu$ parent

Others




FIG. A.7: Other components of $\pi^+ \nu_\mu$ parent

A.2 Flux HP leading to neutrino in NOvA

This appendix contains the hadronic cascade in NuMI leading to a muon neutrino and electron neutrino in the center of the NOvA front face:

- Focusing components (figure A.8).
- Neutrino parents (figure A.9).
- Neutrino grandparents (figure A.10).
- Interaction map (figure A.11 and A.12).
- Hadronic production in the target (figures A.13).
- Beam attenuation (figure A.14).





FIG. A.8: Focusing components for NOvA ND




FIG. A.9: Muon and electron neutrino parents in NOvA







FIG. A.10: Muon and electron neutrino grandparents in NOvA

FIG. A.11: Interaction map for ν_μ in NOvA

FIG. A.12: Incident protons for ν_μ in NOvA

FIG. A.13: Hadron production leading to ν_μ in NOvA




FIG. A.14: Beam attenuation for particles leading to ν_μ in NOvA

APPENDIX B

NuMI flux tables

This appendix contains tables of the Generation 2 flux results:

- MINERvA Gen2-thin Low Energy flux for ν_μ , $\bar{\nu}_\mu$ and ν_e at MINERvA (table B.1).
- MINERvA Gen2-thick Low Energy flux for ν_μ , $\bar{\nu}_\mu$ and ν_e at MINERvA (table B.2).
- MINERvA Gen2-thin Medium Energy flux for ν_μ and ν_e at MINERvA (table B.3).
- NOvA Gen2-thin Low Energy flux for ν_μ at NOvA Near Detector (table B.4).

$\Delta E(\text{GeV})$	ϕ_{ν_μ}	$\delta\phi_{\nu_\mu}^{HP}$	$\delta\phi_{\nu_\mu}^{Foc}$	$\phi_{\bar{\nu}_\mu}$	$\delta\phi_{\bar{\nu}_\mu}^{HP}$	$\delta\phi_{\bar{\nu}_\mu}^{Foc}$	ϕ_{ν_e}	$\delta\phi_{\nu_e}^{HP}$	$\delta\phi_{\nu_e}^{Foc}$
[0, 0.5]	2.82	15.7	2.3	2.17	10.4	2.3	0.08	16.6	2.3
[0.5, 1]	6.75	10.6	3.1	6.35	9.9	3.1	0.19	9.1	3.1
[1, 1.5]	17.36	8.1	2.2	16.35	7.3	2.2	0.34	6.5	2.2
[1.5, 2]	25.86	7.6	2.6	23.53	7.0	2.6	0.41	6.8	2.6
[2, 2.5]	34.00	7.6	2.8	30.32	7.1	2.8	0.44	6.5	2.8
[2.5, 3]	41.73	7.2	3.1	36.84	6.5	3.1	0.39	6.3	3.1
[3, 3.5]	43.79	6.5	3.5	37.39	5.7	3.5	0.36	6.1	3.5
[3.5, 4]	35.09	6.1	5.4	29.00	5.1	5.4	0.32	5.9	5.4
[4, 4.5]	21.18	6.1	9.1	16.81	5.0	9.1	0.26	6.1	9.1
[4.5, 5]	11.54	6.4	10.4	8.75	5.4	10.4	0.23	6.3	10.4
[5, 5.5]	7.05	7.0	7.6	5.15	6.2	7.6	0.18	5.8	7.6
[5.5, 6]	5.02	8.1	4.5	3.46	7.0	4.5	0.15	5.1	4.5
[6, 6.5]	4.08	8.5	3.1	2.62	7.5	3.1	0.11	5.5	3.1
[6.5, 7]	3.41	8.8	2.7	2.14	7.8	2.7	0.09	5.6	2.7
[7, 7.5]	2.97	9.0	2.5	1.77	7.9	2.5	0.06	5.9	2.5
[7.5, 8]	2.62	9.0	2.5	1.48	8.2	2.5	0.06	5.6	2.5
[8, 8.5]	2.34	9.1	2.4	1.24	7.9	2.4	0.04	6.1	2.4
[8.5, 9]	2.11	9.1	2.3	1.09	8.3	2.3	0.04	6.5	2.3
[9, 9.5]	1.91	9.1	2.2	0.99	7.9	2.2	0.04	7.7	2.2
[9.5, 10]	1.74	8.9	2.2	0.88	8.2	2.2	0.04	8.4	2.2
[10, 10.5]	1.58	9.0	2.1	0.74	8.2	2.1	0.03	8.5	2.1
[10.5, 11]	1.47	8.8	2.1	0.71	8.3	2.1	0.03	8.0	2.1
[11, 11.5]	1.33	8.7	2.0	0.62	8.2	2.0	0.03	7.8	2.0
[11.5, 12]	1.21	8.6	2.0	0.58	8.0	2.0	0.02	8.1	2.0
[12, 12.5]	1.10	8.6	2.0	0.50	8.2	2.0	0.02	8.3	2.0
[12.5, 13]	1.01	8.6	2.0	0.47	8.0	2.0	0.02	10.8	2.0
[13, 13.5]	0.91	8.5	2.0	0.42	8.1	2.0	0.02	9.0	2.0
[13.5, 14]	0.86	8.5	2.0	0.37	8.2	2.0	0.02	9.5	2.0
[14, 14.5]	0.78	8.5	2.0	0.34	7.9	2.0	0.02	10.7	2.0
[14.5, 15]	0.71	8.3	2.0	0.31	8.4	2.0	0.02	9.7	2.0
[15, 15.5]	0.64	8.4	2.0	0.27	8.6	2.0	0.01	11.0	2.0
[15.5, 16]	0.59	8.4	2.0	0.26	8.6	2.0	0.01	11.7	2.0
[16, 16.5]	0.55	8.4	2.0	0.23	8.9	2.0	0.02	10.9	2.0
[16.5, 17]	0.49	8.5	2.0	0.19	9.2	2.0	0.01	10.5	2.0
[17, 17.5]	0.45	8.3	2.0	0.19	8.8	2.0	0.01	10.8	2.0
[17.5, 18]	0.41	8.6	2.0	0.17	8.8	2.0	0.01	10.9	2.0
[18, 18.5]	0.37	8.4	2.0	0.15	8.9	2.0	0.01	11.6	2.0
[18.5, 19]	0.34	8.7	2.0	0.13	10.2	2.0	0.01	11.7	2.0
[19, 19.5]	0.32	8.8	2.0	0.12	11.1	2.0	0.01	11.2	2.0
[19.5, 20]	0.29	9.3	2.0	0.12	10.9	2.0	0.01	10.9	2.0

TABLE B.1: Generation2 - thin Low Energy flux results at MINERvA. Flux units: $\nu/m^2/10^6 POT$. Uncertainties are given in percentage.

$\Delta E(GeV)$	ϕ_{ν_μ}	$\delta\phi_{\nu_\mu}^{HP}$	$\delta\phi_{\nu_\mu}^{Foc}$	$\phi_{\bar{\nu}_\mu}$	$\delta\phi_{\bar{\nu}_\mu}^{HP}$	$\delta\phi_{\bar{\nu}_\mu}^{Foc}$	ϕ_{ν_e}	$\delta\phi_{\nu_e}^{HP}$	$\delta\phi_{\nu_e}^{Foc}$
[0, 0.5]	3.54	10.0	2.3	2.88	7.3	2.3	0.08	14.7	2.3
[0.5, 1]	7.35	7.8	3.1	6.86	7.4	3.1	0.19	5.3	3.1
[1, 1.5]	17.29	7.4	2.2	16.31	6.7	2.2	0.33	4.1	2.2
[1.5, 2]	25.36	4.9	2.6	23.74	4.7	2.6	0.41	4.3	2.6
[2, 2.5]	33.72	5.0	2.8	30.96	5.0	2.8	0.44	3.9	2.8
[2.5, 3]	42.91	4.6	3.1	39.03	5.0	3.1	0.38	3.9	3.1
[3, 3.5]	44.38	4.6	3.5	38.46	4.9	3.5	0.36	3.8	3.5
[3.5, 4]	34.54	4.9	5.4	29.25	5.0	5.4	0.32	3.9	5.4
[4, 4.5]	19.75	4.7	9.1	16.92	4.8	9.1	0.26	3.8	9.1
[4.5, 5]	10.36	4.6	10.4	8.69	4.9	10.4	0.23	3.8	10.4
[5, 5.5]	6.16	4.7	7.6	5.02	5.2	7.6	0.17	3.7	7.6
[5.5, 6]	4.32	4.8	4.5	3.36	5.4	4.5	0.14	3.4	4.5
[6, 6.5]	3.46	4.9	3.1	2.51	5.4	3.1	0.10	3.8	3.1
[6.5, 7]	2.85	5.0	2.7	2.01	5.6	2.7	0.08	3.8	2.7
[7, 7.5]	2.46	5.0	2.5	1.63	5.7	2.5	0.06	4.0	2.5
[7.5, 8]	2.16	5.0	2.5	1.35	5.9	2.5	0.06	3.5	2.5
[8, 8.5]	1.95	5.1	2.4	1.12	6.0	2.4	0.04	3.9	2.4
[8.5, 9]	1.77	5.2	2.3	0.99	6.1	2.3	0.04	3.8	2.3
[9, 9.5]	1.61	5.2	2.2	0.91	6.1	2.2	0.03	4.7	2.2
[9.5, 10]	1.47	5.2	2.2	0.82	6.2	2.2	0.03	4.7	2.2
[10, 10.5]	1.34	5.3	2.1	0.70	6.4	2.1	0.03	5.4	2.1
[10.5, 11]	1.25	5.2	2.1	0.66	6.2	2.1	0.03	5.1	2.1
[11, 11.5]	1.13	5.1	2.0	0.58	6.6	2.0	0.02	4.3	2.0
[11.5, 12]	1.04	5.1	2.0	0.54	6.9	2.0	0.02	4.1	2.0
[12, 12.5]	0.96	5.0	2.0	0.47	6.8	2.0	0.02	4.9	2.0
[12.5, 13]	0.90	5.0	2.0	0.44	6.7	2.0	0.02	6.4	2.0
[13, 13.5]	0.83	4.8	2.0	0.40	6.5	2.0	0.02	4.7	2.0
[13.5, 14]	0.80	4.9	2.0	0.36	6.5	2.0	0.02	5.5	2.0
[14, 14.5]	0.75	4.9	2.0	0.33	6.7	2.0	0.01	4.4	2.0
[14.5, 15]	0.69	4.9	2.0	0.30	7.0	2.0	0.01	4.1	2.0
[15, 15.5]	0.63	4.8	2.0	0.26	7.0	2.0	0.01	5.2	2.0
[15.5, 16]	0.59	4.7	2.0	0.24	7.3	2.0	0.01	5.4	2.0
[16, 16.5]	0.55	4.7	2.0	0.21	7.6	2.0	0.01	4.7	2.0
[16.5, 17]	0.50	4.7	2.0	0.18	8.0	2.0	0.01	5.3	2.0
[17, 17.5]	0.46	4.7	2.0	0.17	8.2	2.0	0.01	4.9	2.0
[17.5, 18]	0.42	4.7	2.0	0.15	7.8	2.0	0.01	4.7	2.0
[18, 18.5]	0.38	4.9	2.0	0.13	7.7	2.0	0.01	5.0	2.0
[18.5, 19]	0.34	5.0	2.0	0.12	8.4	2.0	0.01	5.7	2.0
[19, 19.5]	0.32	5.2	2.0	0.11	8.8	2.0	0.01	5.7	2.0
[19.5, 20]	0.30	5.6	2.0	0.10	9.0	2.0	0.01	5.1	2.0

TABLE B.2: Generation2 - thick Low Energy flux results at MINERvA. Flux units: $\nu/m^2/10^6 POT$. Uncertainties are given in percentage.

ΔE	ϕ_{ν_μ}	$\delta\phi_{\nu_\mu}^{HP}$	$\delta\phi_{\nu_\mu}^{Foc}$	$\phi_{\bar{\nu}_\mu}$	$\delta\phi_{\bar{\nu}_\mu}^{HP}$	ϕ_{ν_e}	$\delta\phi_{\nu_e}^{HP}$	$\phi_{\bar{\nu}_e}$	$\delta\phi_{\bar{\nu}_e}^{HP}$
[0, 0.5]	2.45	16.5	12.8	1.64	15.5	0.04	17.3	0.00	79.4
[0.5, 1]	4.30	12.6	8.4	1.61	16.2	0.13	8.7	0.01	19.1
[1, 1.5]	4.07	11.9	5.2	1.38	15.3	0.20	8.2	0.04	18.9
[1.5, 2]	8.39	9.2	3.6	1.32	15.6	0.29	7.5	0.02	17.4
[2, 2.5]	19.96	8.1	3.0	1.22	14.0	0.36	7.0	0.02	16.0
[2.5, 3]	33.83	7.5	2.3	1.27	13.1	0.41	7.2	0.02	14.0
[3, 3.5]	44.07	7.3	3.1	1.17	12.1	0.44	6.6	0.03	13.5
[3.5, 4]	48.68	7.4	3.8	1.24	11.5	0.45	6.9	0.02	17.5
[4, 4.5]	53.37	7.7	3.4	1.23	10.9	0.44	6.8	0.01	13.9
[4.5, 5]	60.55	7.9	3.2	1.16	10.8	0.42	7.1	0.01	11.2
[5, 5.5]	66.98	7.8	3.2	1.15	10.4	0.40	7.1	0.01	14.6
[5.5, 6]	69.57	7.5	3.5	1.08	9.9	0.38	6.8	0.02	13.1
[6, 6.5]	66.23	7.1	4.6	1.01	9.9	0.34	6.5	0.01	9.9
[6.5, 7]	56.85	6.8	6.7	0.96	9.1	0.31	6.5	0.01	10.5
[7, 7.5]	43.61	6.7	9.4	0.84	9.3	0.27	6.5	0.01	10.5
[7.5, 8]	30.81	6.6	12.5	0.77	9.5	0.23	7.3	0.01	13.0
[8, 8.5]	20.79	6.7	15.7	0.65	8.7	0.20	6.7	0.01	12.7
[8.5, 9]	13.53	6.8	18.4	0.58	8.6	0.18	6.3	0.01	9.8
[9, 9.5]	8.75	7.0	19.4	0.54	8.7	0.15	6.6	0.01	13.7
[9.5, 10]	5.76	7.3	18.1	0.49	8.8	0.14	6.3	0.01	11.4
[10, 10.5]	4.03	7.6	15.9	0.46	8.7	0.12	6.4	0.01	12.0
[10.5, 11]	2.96	8.1	13.4	0.42	9.1	0.10	7.0	0.01	11.5
[11, 11.5]	2.30	8.4	11.6	0.38	9.0	0.10	7.0	0.01	11.8
[11.5, 12]	1.92	8.5	8.0	0.36	8.9	0.08	6.3	0.00	16.7
[12, 12.5]	1.64	8.7	5.7	0.33	9.7	0.07	6.4	0.01	10.9
[12.5, 13]	1.40	8.7	6.0	0.29	9.3	0.06	6.9	0.01	16.1
[13, 13.5]	1.29	8.6	3.1	0.27	9.4	0.06	7.0	0.01	13.1
[13.5, 14]	1.16	8.4	2.3	0.25	9.8	0.05	7.0	0.01	9.5
[14, 14.5]	1.03	8.4	2.1	0.22	10.1	0.04	6.9	0.00	12.4
[14.5, 15]	0.93	8.1	3.0	0.19	9.5	0.04	7.4	0.00	14.9
[15, 15.5]	0.86	8.3	4.4	0.19	10.3	0.03	7.0	0.01	12.4
[15.5, 16]	0.78	8.3	2.7	0.16	10.3	0.03	8.4	0.00	14.9
[16, 16.5]	0.73	7.9	2.4	0.14	10.7	0.03	8.9	0.00	10.3
[16.5, 17]	0.67	8.2	3.8	0.13	10.9	0.03	8.6	0.01	12.9
[17, 17.5]	0.62	8.1	4.2	0.12	10.7	0.02	9.6	0.00	10.6
[17.5, 18]	0.56	8.1	5.7	0.10	12.2	0.03	9.3	0.00	13.5
[18, 18.5]	0.54	7.6	4.9	0.09	11.1	0.02	10.6	0.00	14.4
[18.5, 19]	0.50	7.6	3.9	0.08	11.8	0.02	9.7	0.00	18.6
[19, 19.5]	0.48	7.6	4.4	0.07	13.3	0.02	9.5	0.00	12.9
[19.5, 20]	0.45	7.5	2.9	0.07	14.2	0.01	10.2	0.00	11.1

TABLE B.3: Generation2 - thin Medium Energy flux results at MINERvA. Flux units: $\nu/m^2/10^6 POT$. Uncertainties are given in percentage.

$\Delta E(\text{GeV})$	ϕ_{ν_μ}	$\delta\phi_{\nu_\mu}^{HP}$	$\Delta E(\text{GeV})$	ϕ_{ν_μ}	$\delta\phi_{\nu_\mu}^{HP}$
[0, 0.1]	0.25	31.3	[0.1, 0.2]	0.62	23.0
[0.2, 0.3]	0.82	16.7	[0.3, 0.4]	1.00	13.3
[0.4, 0.5]	1.19	11.7	[0.5, 0.6]	1.38	10.9
[0.6, 0.7]	1.56	10.4	[0.7, 0.8]	1.70	10.0
[0.8, 0.9]	1.84	9.7	[0.9, 1]	2.03	9.3
[1, 1.1]	2.30	9.1	[1.1, 1.2]	2.71	8.8
[1.2, 1.3]	3.36	8.5	[1.3, 1.4]	4.26	8.3
[1.4, 1.5]	5.14	8.1	[1.5, 1.6]	5.93	8.0
[1.6, 1.7]	6.63	7.9	[1.7, 1.8]	7.18	7.9
[1.8, 1.9]	7.47	7.8	[1.9, 2]	7.43	7.7
[2, 2.1]	7.02	7.7	[2.1, 2.2]	6.29	7.7
[2.2, 2.3]	5.38	7.6	[2.3, 2.4]	4.44	7.5
[2.4, 2.5]	3.59	7.5	[2.5, 2.6]	2.87	7.4
[2.6, 2.7]	2.31	7.3	[2.7, 2.8]	1.85	7.2
[2.8, 2.9]	1.47	7.3	[2.9, 3]	1.17	7.2
[3, 3.1]	0.93	7.4	[3.1, 3.2]	0.73	7.5
[3.2, 3.3]	0.58	8.1	[3.3, 3.4]	0.46	8.4
[3.4, 3.5]	0.37	8.7	[3.5, 3.6]	0.32	9.2
[3.6, 3.7]	0.28	9.8	[3.7, 3.8]	0.25	10.0
[3.8, 3.9]	0.23	10.0	[3.9, 4]	0.21	10.0
[4, 4.1]	0.20	10.6	[4.1, 4.2]	0.18	10.6
[4.2, 4.3]	0.18	10.7	[4.3, 4.4]	0.16	10.7
[4.4, 4.5]	0.15	10.5	[4.5, 4.6]	0.15	10.7
[4.6, 4.7]	0.14	10.2	[4.7, 4.8]	0.14	10.0
[4.8, 4.9]	0.13	10.3	[4.9, 5]	0.13	10.8
[5, 5.1]	0.12	10.3	[5.1, 5.2]	0.11	10.7
[5.2, 5.3]	0.11	10.0	[5.3, 5.4]	0.11	10.5
[5.4, 5.5]	0.10	10.4	[5.5, 5.6]	0.10	10.4
[5.6, 5.7]	0.09	9.8	[5.7, 5.8]	0.09	10.2
[5.8, 5.9]	0.09	10.1	[5.9, 6]	0.09	9.8

TABLE B.4: Generation2 - thin Medium Energy flux results at the center of the front face of NOvA Near Detector. Flux units: $\nu/m^2/10^6 POT$. Uncertainties are given in percentage.

BIBLIOGRAPHY

- [1] P. Adamson et al. (MINOS Collaboration), Phys. Rev. D **77**, 072002 (2008).
- [2] Dune Collaboration, *Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE) Conceptual Design Report*, FERMILAB-DESIGN-2016-02.
- [3] J. A. Formaggio and G. P. Zeller, Rev. Mod. Phys. **84**, 1307 (2012).
- [4] MINERvA Collaboration, *Proposal to Perform a High-Statistics Neutrino Scattering Experiment using a Fine grained Detector in the NuMI Beam* (Batavia IL, 2004).
- [5] A. Higuera et al. (MINERvA Collaboration), Phys. Rev. Lett. **113**, 261802 (2014).
- [6] Sacha E. Kopp, Phys.Rept **439**, 101 (2007).
- [7] van der Meer, S. and Vahlbruch, K. M., CERN-1963-037 pp. 97–110 (1963).
- [8] P. Adamson et. al., Nucl.Instrum.Meth. **A806**, 279 (2016).
- [9] NuMI-MINOS, *NuMI technical design handbook*, Accessed: 2016-01-25, URL http://www-numi.fnal.gov/numwork/tdh/tdh_index.html.
- [10] Fermilab, *Accelerator report no. 4: The linac*, Accessed: 2016-01-25, URL http://www.fnal.gov/pub/news04/update_archive/update_10-1.html.

- [11] Fermilab, *Booster department*, Accessed: 2016-02-02, URL <http://www-ad.fnl.gov/proton/booster.html>.
- [12] C. C. Jensen, G. A. Krafczyk (Fermilab), Conf.Proc. **C0505161**, 692 (2005).
- [13] K. Budal, IEEE Trans. Nucl. Sci. **14**, 1132 (1967).
- [14] Fermilab, *Visual media service*, Accessed: 2016-02-22, URL <http://vms.fnal.gov>.
- [15] GEANT4 Collaboration, Nuclear Instruments and Methods in Physics Research A **506**, 250 (2003).
- [16] Geant4, *Model: Ftfp*, Accessed: 2016-02-11, URL http://geant4.cern.ch/support/proc_mod_catalog/models/hadronic/FTFP.html.
- [17] Geant4, *Model: Bertini cascade*, Accessed: 2016-02-11, URL http://geant4.cern.ch/support/proc_mod_catalog/models/hadronic/BertiniCascade.html.
- [18] Andreopoulos, C. et al, Nuclear Instruments and Methods in Physics Research A **614**, 87 (2010).
- [19] M. Colaboration, *The minos technical design report*, Accessed: 2016-03-01, URL http://www-numi.fnal.gov/minwork/info/minos_tdr.html.
- [20] Ž. Pavlović, Ph.D. thesis (2008), Fermilab-Thesis-2008-59.
- [21] M. Kordosky, *Reassessment of the skin depth effect* (2014).
- [22] P. L. Brun, Tech. Rep. (2014).
- [23] J. Hylen, *Target location along beam-line; summary of entire minos run* (2014).
- [24] P. L. Brun, Tech. Rep. (2014).

- [25] L. J. Loiacono, Ph.D. thesis (2011), FERMILAB-THESIS-2011-06.
- [26] M. Apollonio et al. (HARP Collaboration), Phys. Rev. C **80**, 035208 (2009).
- [27] M. Apollonio et al. (HARP Collaboration), Nuclear Physics A **821**, 112 (2009).
- [28] M. Apollonio et al. (HARP Collaboration), Phys. Rev. C **82**, 045208 (2010).
- [29] M. Apollonio et al. (HARP Collaboration), Eur. Phys. J. C **52**, 29 (2007).
- [30] D. W. Schmitz, Ph.D. thesis (2008), FERMILAB-THESIS-2008-26.
- [31] D. A. A. Wickremasinghe, Ph.D. thesis (2015), FERMILAB-THESIS-2015-23.
- [32] Fermilab, *Booster neutrino experiment*, Accessed: 2016-03-8, URL <https://www-boone.fnal.gov/>.
- [33] A. A. Aguilar-Arevalo et al. (MiniBooNE Collaboration), Phys. Rev. D **79**, 072002 (2009).
- [34] K. Abe et al. (T2K Collaboration), Phys. Rev. D **87**, 012001 (2013).
- [35] N. abgrall et al. (NA61/SHINE Collaboration), Phys. Rev. C **84**, 034604 (2011).
- [36] N. abgrall et al. (NA61/SHINE Collaboration), Phys. Rev. C **85**, 035210 (2012).
- [37] N. Abgrall et al. (NA61/SHINE Collaboration), Nucl. Instrum. Meth. **A701**, 99 (2013).
- [38] Bonesini, M. et al., Eur. Phys. J. **C20**, 13 (2001).
- [39] P. Adamson et al. (MINOS Collaboration), Phys. Rev. D **81**, 072002 (2010).
- [40] A. Bodek, U. Sarica, D. Naples, L. Ren, The European Physical Journal C **72-4**, 1973 (2012).

- [41] W. G. Seligman, Ph.D. thesis (1997), FERMILAB-THESIS-1997-21.
- [42] D. Bhattacharya, Ph.D. thesis (2009), FERMILAB-THESIS-2009-11.
- [43] J. Devan, Ph.D. thesis (2015), Fermilab-Thesis-2015-29.
- [44] J. Park, Ph.D. thesis (2013), FERMILAB-THESIS-2013-36.
- [45] Richard P. Feynman, Phys. Rev. Lett. **23**, 1415 (1969).
- [46] Benecke J., Chou T.T., Yang C.N and Yen E., Phys. Rev. **188**, 2159 (1969).
- [47] T.T. Böhlen, F. Cerutti, M.P.W. Chin, A. Fassò, A. Ferrari, P.G. Ortega, A. Mairani, P.R. Sala, G. Smirnov and V. Vlachoudis, Nuclear Data Sheets **120**, 211 (2014).
- [48] A. Ferrari, P.R. Sala, A. Fasso, and J. Ranft, CERN-2005-10, INFN/TC_05/11, SLAC-R-773 (2005).
- [49] M. Kordosky, *Energy dependence of hadron production*.
- [50] J. Wolcott, Tech. Rep. (2013).
- [51] D. S. Barton et al., Phys. Rev. D **27**, 2580 (1983).
- [52] P. Skubic, et al., Phys. Rev. D **18**, 3115 (1978).
- [53] T. Golan, *Uncertainty coming from a -scaling*.
- [54] M. Jerkins, Tech. Rep. (2012).
- [55] M. Kordosky, Tech. Rep. (2012).
- [56] W.H.Press et Al., *Numerical Recipes. The Art of Scientific Computing* (Cambridge, University Press, 2007), 3rd ed.

- [57] J. Paley et Al. (MIPP Collaboration), Phys. Rev. D **90**, 032001 (2014).
- [58] S. M. Seun, Ph.D. thesis (2007), FERMILAB-THESIS-2007-61.
- [59] G. Bellettini et al., Nucl. Phys. **79**, 609 (1966).
- [60] S. P. Denisov et al., Nucl. Phys. **B61**, 62 (1973).
- [61] NA49 Collaboration, Eur.Phys.J. **C49**, 897 (2007).
- [62] NA49 Collaboration, Eur.Phys.J. **C73**, 2364 (2013).
- [63] G. M. Tinti, Ph.D. thesis (2010), FERMILAB-THESIS-2010-44.
- [64] A. V. Lebedev, Ph.D. thesis (2007), FERMILAB-THESIS-2007-76.
- [65] NA49 Collaboration, Nucl.Instrum.Meth. **A430**, 210 (1999).
- [66] CERN, *Large acceptance hadron detector for an investigation of pb-induced reactions at the cern sps*, Accessed: 2016-03-06, URL <http://na49info.web.cern.ch>.
- [67] Antonino Zichichi, *Understanding the Fundamental Constituents of Matter* (Plenum Press, 1978).
- [68] A. S. Carroll et al., Physics Letter **B**, 319 (1979).
- [69] T. J. Roberts et al., Nuclear Physics **B159**, 56 (1979).
- [70] J. W. Cronin et al., Phys.Rev. **107**, 1121 (1957).
- [71] J. V. Allaby et al., Yad.Fiz. **12**, 538 (1970).
- [72] B. W. Allardyce et al., Nuclear Physics **209**, 1 (1973).
- [73] M. Longo et al., Phys. Rev. **125**, 701 (1962).

- [74] V. B. Fedorov et al., Sov.J.Nucl.Phys. **222**, 413 (1978).
- [75] B. M. Bobchenko et al., Sov.J.Nucl.Phys. **30**, 805 (1979).
- [76] R. J. Abrams et al., Phys.Rev. **D1**, 1917 (1970).
- [77] R. Hatcher, *Proposal for a unified flux n-tuple format* (2014).
- [78] L. Aliaga, *Flux hadron production reweighting (coll. meeting march 2015)*.
- [79] L. Aliaga, *Flux reweighting update (coll. meeting june 2015)*.
- [80] L. Aliaga, *Ppfx update (coll. meeting september 2015)*.
- [81] G. A. Fiorentini et al. (MINERvA Collaboration), Phys. Rev. Lett. **111**, 022502 (2013).
- [82] L. Fields et al. (MINERvA Collaboration), Phys. Rev. Lett. **111**, 022501 (2013).
- [83] B.G. Tice et al. (MINERvA Collaboration), Phys. Rev. Lett. **112**, 231801 (2014).
- [84] T. Walton et al. (MINERvA Collaboration), Phys. Rev. D. **91**, 071301 (2015).
- [85] T. Le et al. (MINERvA Collaboration), Phys. Lett. B. **749**, 130 (2015).
- [86] B. Eberly et al. (MINERvA Collaboration), Phys. Lett. D. **92**, 092008 (2015).
- [87] M. Kordosky, *Comparisons of the gen-2 and low-v fluxes* (2016).
- [88] J. Park, *Neutrino electron scattering update*.
- [89] L. Fields, *Neutrino electron scattering plot approval*.
- [90] Jeremy Wolcott et al. (MINERvA Collaboration), Phys. Rev. D **81**, 072002 (2010).

- [91] A. R. for the NuMIX working group, *A document describing how the on and off axis near detectors for nova and minos sample the numi beam during a variety of beam modes.*